

QUANTITATIVE HOMOGENIZATION OF DEGENERATE RANDOM ENVIRONMENTS

ARIANNA GIUNTI, JEAN-CHRISTOPHE MOURRAT

ABSTRACT. We study discrete linear divergence-form operators with random coefficients, also known as the random conductance model. We assume that the conductances are bounded, independent and stationary; the law of a conductance may depend on the orientation of the associated edge. We give a simple necessary and sufficient condition for the relaxation of the environment seen by the particle to be diffusive, in the sense of every polynomial moment. As a consequence, we derive polynomial moment estimates on the corrector.

MSC 2010: 35B27, 35K65, 60K37.

KEYWORDS: Quantitative homogenization, environment viewed by the particle, mixing of Markov chains, corrector estimate.

1. INTRODUCTION

We study the homogenization of discrete divergence-form operators

$$(1.1) \quad -\nabla \cdot a \nabla f(x) := \sum_{y \sim x} a((x, y))(f(y) - f(x)),$$

where $a = (a(e))_{e \in \mathbb{E}}$ is a family of independent random variables indexed by the nearest-neighbor, unoriented edges of the graph $\mathbb{Z}^d$, $d \geq 2$. We assume that the coefficients $a(e)$ take values in $[0, 1]$, that the law of $a(e)$ depends only on the orientation of the edge e , and that none of these d probability laws is a Dirac mass at 0. In this setting, we give a necessary and sufficient condition for the relaxation of the “environment viewed by the particle” to be diffusive in the sense of every polynomial moment. As the name implies, this process can be described in terms of the random walk with generator given by (1.1). We will rather define the flow of its semigroup directly by means of the PDE (1.6) below. Denoting by u_t the solution to (1.6) with bounded, local and centered initial condition g , we show that the property

$$(1.2) \quad \text{for every } p \geq 1, \quad \sup_{t \geq 1} t^{\frac{d}{2}} \langle |u_t|^{2p} \rangle^{\frac{1}{p}} < \infty$$

holds as soon as

$$(1.3) \quad \text{for every } q \geq 1, \quad \langle \left(\sup_{1 \leq i \leq d} a(e_i) \right)^{-q} \rangle < \infty,$$

where $(e_1, \dots, e_d)$ is the canonical basis of $\mathbb{Z}^d$, and $\langle \cdot \rangle$ denotes the expectation with respect to the law of $\{a(e)\}$. Moreover, if (1.2) holds for a single “generic” bounded and local initial datum g , then (1.3) holds as well. Finally, we show that if the initial condition g in (1.6) is the divergence of a local bounded function and if (1.3) holds, then (1.2) can be improved to

$$(1.4) \quad \text{for every } p \geq 1 \text{ and } \varepsilon > 0, \quad \sup_{t \geq 1} t^{\frac{d}{2}+1-\varepsilon} \langle |u_t|^{2p} \rangle^{\frac{1}{p}} < \infty.$$

A rather degenerate example of environment satisfying (1.3) can be constructed by letting $a(e)$ be i.i.d. Bernoulli random variables in $(d-1)$ directions, and letting

$a(e) = (1 + E_e)^{-1}$ with (E_e) i.i.d. exponential random variables in the remaining direction.

As shown in [9, 5] and recalled below, these estimates imply a range of other quantitative homogenization results, including bounds on the corrector. They can also be used to prove quenched central limit theorems for the associated random walk.

Under the condition that the coefficients $a(e)$ are uniformly bounded away from 0 and infinity, the estimate (1.2) and the stronger estimate (1.4) with $\varepsilon = 0$ were proved in [5]. We refer to [3, 6, 7, 2] for more recent developments under this assumption. Bounds on the corrector under assumptions similar to ours and for $d \geq 3$ were obtained in [8]. The case of coefficients that are bounded away from 0 but not from infinity was considered in [9, 4].

Our approach is inspired by the strategy of [5], which rests on quenched heat kernel bounds. In the context of degenerate environments, quenched diffusive bounds on the heat kernel are false in general. However, under the condition (1.3), we will be able to control the anomalous behavior of the heat kernel, in the sense of every polynomial moment, by exploiting the method presented in [10]. We then show that these weaker bounds are sufficient to imply (1.2) and (1.4).

Our long-term goal is to develop a comparable strategy in the context of interacting particle systems, in particular to study the relaxation of the environment viewed by a tagged particle in the symmetric exclusion process. We expect the results of this paper to be a first step in this direction. Loosely speaking, in the present work, we leverage on the existence of one “good direction” where conductances are well-behaved. For the exclusion process, we hope to benefit from the good behavior of the model in the *time* direction, as illustrated for instance by [10, Lemma 5.3].

1.1. Notation and main result. We say that $x, y \in \mathbb{Z}^d$ are neighbors, and write $x \sim y$, when $|x - y| = 1$. This endows $\mathbb{Z}^d$ with a graph structure, so that we may introduce the associated set of unoriented edges $\mathbb{B}$. Throughout the paper, we will typically denote points of $\mathbb{Z}^d$ by x, y, z , and edges in $\mathbb{B}$ by b, e . For a given edge $e \in \mathbb{B}$, we write $\underline{e}$ and $\bar{e}$ to denote its two endpoints, with the convention that $\bar{e} - \underline{e} = e_i$ for some $i \in \{1, \dots, d\}$, where we recall that $(e_1, \dots, e_d)$ is the canonical basis of $\mathbb{Z}^d$. We identify the vector $e_i \in \mathbb{Z}^d$ with the edge $(0, e_i)$, for each $i \in \{1, \dots, d\}$.

The space of “environments” we consider is $\Omega := [0, 1]^{\mathbb{B}}$. The group $\mathbb{Z}^d$ naturally acts by translations on Ω in the following way: for every $x \in \mathbb{Z}^d$ and $a = (a(e))_{e \in \mathbb{B}} \in \Omega$, we define

$$(1.5) \quad \tau_x a := (a(x + e))_{e \in \mathbb{B}},$$

where for $e \in \mathbb{B}$, we write $x + e := (x + \underline{e}, x + \bar{e})$. We consider a random $a = (a(e))_{e \in \mathbb{B}} \in \Omega$ whose law we denote by $\langle \cdot \rangle$. We assume the family of random variables $a = (a(e))$ to be independent and stationary, i.e. for every $x \in \mathbb{Z}^d$, the random variables $\tau_x a$ and a have the same law. In other words, the random variables $a(e)$ are independent, and the law of $a(e)$ only depends on the orientation of the edge e . We assume that for every e , $\mathbb{P}[a(e) = 0] < 1$.

For a random variable $\xi : \Omega \rightarrow \mathbb{R}$ and a fixed edge $b \in \mathbb{B}$, we define

$$D_b \xi = D_b \xi(a) := \xi(\tau_{\bar{b}} a) - \xi(\tau_{\underline{b}} a),$$

and simply write $D\xi$ for the d -dimensional random vector defined as

$$(D\xi)_i := D_{e_i} \xi.$$

We observe that for every $p \in [1, +\infty]$ the operator $D : L^p(\Omega) \rightarrow L^p(\Omega)^d$ is bounded and that its adjoint in $L^2(\Omega)$, which we denote by $D^* : L^2(\Omega)^d \rightarrow L^2(\Omega)$, is defined

as

$$D^* \xi := \sum_{i=1}^d D_i^* \xi_i(a), \quad \text{where} \quad D_i^* \xi_i := \xi(\tau_{-e_i} a) - \xi(a).$$

Given a random variable $g \in L^1(\Omega)$, with $\langle g \rangle = 0$, our goal is to understand the relaxation to equilibrium of the solution $u : \mathbb{R}_+ \times \Omega \rightarrow \mathbb{R}$ to

$$(1.6) \quad \begin{cases} \partial_t u_t + D^* a D u_t = 0 \\ u_0 = g, \end{cases}$$

where

$$D^* a D u_t(a) := \sum_{i=1}^d D_i^*(a(e_i) D_i u_t(a)) = \sum_{e \ni 0} a(e) D_e u_t(a).$$

Whenever no confusion occurs, we write u_t instead of $u_t(a)$. We say that a function $g : \Omega \rightarrow \mathbb{R}$ is local if $g(a)$ depends only on a finite number of conductances $a(e^{(1)}), \dots, a(e^{(N)})$. In this case, we say that g has support of size N . Here is our main result.

Theorem 1.1. *Under the moment condition (1.3), the following statements hold.*

- (a) *For every $p \in [1, \infty)$, there exists a constant $C = C(d, p) < +\infty$ such that if $g : \Omega \rightarrow \mathbb{R}$ is local with support of size N , bounded and centered, then*

$$(1.7) \quad \langle |u_t|^{2p} \rangle^{1/p} \leq CN \|g\|_{L^\infty(\Omega)}^2 t^{-\frac{d}{2}}.$$

- (b) *For every $p \in [1, \infty)$ and $\varepsilon > 0$, there exists a constant $C = C(d, p, \varepsilon) < \infty$ such that if $f : \Omega \rightarrow \mathbb{R}$ is local with support of size N and bounded, and if $g = D^* f$, then*

$$(1.8) \quad \langle |u_t|^{2p} \rangle^{1/p} \leq CN \|f\|_{L^\infty(\Omega)}^2 t^{-(\frac{d}{2} + 1 - \varepsilon)}.$$

1.2. Consequences of the main result. As was shown in [9, Section 9] and [5, Section 6], Theorem 1.1 implies a host of other results of interest in stochastic homogenization. In particular, estimates on the corrector can be derived from the relaxation to equilibrium of the solution to (1.6) with $g = D^*(ae)$, by integration in time.

Corollary 1.2. *Assume that the moment condition (1.3) holds, and let $e \in \mathbb{B}$. If $d > 2$, then there exists $\phi_e \in L^2(\Omega)$ solution to the equation*

$$(1.9) \quad D^* a D \phi_e = -D^* a e,$$

and moreover, ϕ_e is in $L^p(\Omega)$ for every $p \in [1, \infty)$. If $d = 2$, then there exists χ in the $L^2(\Omega)^d$ -closure of $\{D\phi : \phi \in L^2(\Omega)\}$ solution to the equation

$$(1.10) \quad D^* a \chi = -D^* a e$$

such that $\chi \in L^p(\Omega)^d$ for every $p \in [1, \infty)$.

We refer to [5, Propositions 4 and 6] for proofs of these results. As another example, we can estimate the corrector $\phi_{e,\mu}$ with massive term $\mu > 0$, i.e. the solution of

$$\mu \phi_{e,\mu} + D^* a D \phi_{e,\mu} = -D^* a e.$$

By [5, Proposition 5], we obtain that for every $\varepsilon > 0$ and $p \in [1, \infty)$,

$$\begin{aligned} \sup_{\mu > 0} \mu^\varepsilon \langle |\phi_{e,\mu}|^p \rangle^{\frac{1}{p}} &< \infty & \text{if } d = 2, \\ \sup_{\mu > 0} \langle |\phi_{e,\mu}|^p \rangle^{\frac{1}{p}} &< \infty & \text{if } d > 2. \end{aligned}$$

1.3. On the necessity of the moment condition. We now explain why our assumption (1.3) on the law of $(a(e))$ is necessary in order to have the optimal relaxation decay (1.2). In this subsection we introduce the notation $\lesssim$ for $\leq C$ where the constant C only depends on the dimension d of the lattice $\mathbb{Z}^d$.

If (1.3) does not hold, then we can find $p_0 \geq 1$ and a sequence $\{\varepsilon_{p_0, n}\}_{n \in \mathbb{N}}$, $\varepsilon_{p_0, n} \downarrow 0$ such that for every $n \in \mathbb{N}$,

$$(1.11) \quad \mathbb{P}\left(\sup_{i=1, \dots, d} a(e_i) \leq \varepsilon_{p_0, n}\right) = \prod_{i=1}^d \mathbb{P}_i(a(e_i) \leq \varepsilon_{p_0, n}) \geq \varepsilon_{p_0, n}^{p_0}.$$

We show that the solution u_t of

$$(1.12) \quad \begin{cases} \partial_t u_t + D^* a D u_t = 0 \\ u_0 = g, \end{cases}$$

with $g = a(\tilde{e}) - \langle a(\tilde{e}) \rangle$, for a fixed $\tilde{e}$ such that $0 \notin \tilde{e}$, does not satisfy the bound (1.2). We make this choice of initial datum g for convenience, but as will be seen shortly, this is inessential. From (1.6), we may bound by stationarity

$$|\partial_t u_t| \lesssim \left[\sup_{e \geq 0} a(e) \right] \|u_t\|_{L^\infty} = \left[\sup_{i=1, \dots, d} a(\pm e_i) \right] \|u_t\|_{L^\infty},$$

and hence, by the maximum principle,

$$|\partial_t u_t| \lesssim \left[\sup_{i=1, \dots, d} a(\pm e_i) \right] \|g\|_{L^\infty}.$$

Therefore, if $\sup_{i=1, \dots, d} a(\pm e_i) \leq \frac{1}{t^2}$, we get

$$(1.13) \quad |u_t| \geq |g| - \frac{\|g\|_{L^\infty(\Omega)}}{t},$$

and thus

$$\begin{aligned} \langle |u_t|^{2q} \rangle &\geq \langle |u_t|^{2q} \mathbf{1}_{\sup_i a(\pm e_i) \leq \frac{1}{t^2}} \rangle = \langle |u_t|^{2q} \mid \sup_i a(\pm e_i) \leq \frac{1}{t^2} \rangle \mathbb{P}(\sup_i a(\pm e_i) \leq \frac{1}{t^2}) \\ &\stackrel{(1.13)}{\gtrsim} \langle \left(|g| - \frac{\|g\|_{L^\infty(\Omega)}}{t} \right)^{2q} \mid \sup_i a(\pm e_i) \leq \frac{1}{t^2} \rangle \mathbb{P}(\sup_i a(\pm e_i) \leq \frac{1}{t^2}) \\ &\stackrel{a(\tilde{e}) \perp a(\pm e_i)}{=} \langle \left(|g| - \frac{\|g\|_{L^\infty(\Omega)}}{t} \right)^{2q} \rangle \mathbb{P}(\sup_i a(\pm e_i) \leq \frac{1}{t^2}) \\ &= \langle \left(|g| - \frac{\|g\|_{L^\infty(\Omega)}}{t} \right)^{2q} \rangle \mathbb{P}(\sup_i a(e_i) \leq \frac{1}{t^2}, \sup_i a(-e_i) \leq \frac{1}{t^2}) \\ (1.14) \quad &= \langle \left(|g| - \frac{\|g\|_{L^\infty(\Omega)}}{t} \right)^{2q} \rangle [\mathbb{P}(\sup_i a(e_i) \leq \frac{1}{t^2})]^2, \end{aligned}$$

where in the last equality we use that $\sup_i a(e_i)$ and $\sup_i a(-e_i)$ are independent and have the same law. For $\{t_n\}$ with $t_n^2 = \frac{1}{\varepsilon_{p_0, n}}$ we may apply (1.11) and estimate

$$\langle |u_{t_n}|^{2q} \rangle^{\frac{1}{q}} \gtrsim \langle \left(|g| - \frac{\|g\|_{L^\infty(\Omega)}}{t_n} \right)^{2q} \rangle^{\frac{1}{q}} t_n^{-4 \frac{p_0}{q}} \gtrsim (\langle |g|^{2q} \rangle^{\frac{1}{q}} - \frac{\|g\|_{L^\infty(\Omega)}^2}{t_n^2}) t_n^{-4 \frac{p_0}{q}}.$$

Thus, for any $q > \frac{8p_0}{d}$ we contradict (1.2).

1.4. Organisation of the paper. In the rest of the paper, we assume that the moment condition (1.3) holds. We derive the necessary heat kernel bounds in Section 2, and proceed to prove Theorem 1.1 in Section 3.

2. HEAT KERNEL BOUNDS

We say that a random field $\zeta : \Omega \times \mathbb{Z}^d \rightarrow \mathbb{R}$ is stationary if for every $x \in \mathbb{Z}^d$, we have $\zeta(a, x) = \zeta(\tau_x a, 0)$. Conversely, given a random variable $\xi : \Omega \rightarrow \mathbb{R}$, we define its stationary extension $\bar{\xi} : \Omega \times \mathbb{Z}^d \rightarrow \mathbb{R}$ as the random field given by $\bar{\xi}(a, x) := \xi(\tau_x a)$. If the function $u : \mathbb{R}_+ \times \Omega \rightarrow \mathbb{R}$ solves (1.6), then its stationary extension $\bar{u}_t(x, a) = u_t(\tau_x a)$ is a solution in $\mathbb{R}_+ \times \mathbb{Z}^d$ of the parabolic PDE

$$(2.15) \quad \begin{cases} \partial_t \bar{u}_t - \nabla \cdot \bar{a} \nabla \bar{u}_t = 0 \\ \bar{u}_0 = \bar{g}, \end{cases}$$

with ∇ the spacial discrete gradient defined for an edge $b \in \mathbb{B}$ and a random field ζ as

$$\begin{aligned} \nabla \zeta(a, b) &:= \zeta(a, \bar{b}) - \zeta(a, \underline{b}), \\ a \nabla \zeta(a, b) &:= a(b) \nabla \zeta(a, b). \end{aligned}$$

Let $p_t = p_t(a, x, y)$ be the parabolic Green function associated to the operator $\partial_t - \nabla \cdot \bar{a} \nabla$, i.e. for every $y \in \mathbb{Z}^d$ the unique bounded solution in $\mathbb{Z}^d$ of

$$(2.16) \quad \begin{cases} \partial_t p_t - \nabla \cdot \bar{a} \nabla p_t = 0 \\ p_0(x, y) = \mathbf{1}_{\{x=y\}}(x). \end{cases}$$

For every $\alpha \in \mathbb{R}$, $x \in \mathbb{R}^d$ and $t \geq 0$, we write

$$(2.17) \quad \omega_\alpha(x, t) := \left(\frac{|x|^2}{t+1} + 1 \right)^{\frac{\alpha}{2}}.$$

The goal of this section is to show the heat kernel upper bound summarized in Lemma 2.1, which we then lift to an estimate on the gradient of the heat kernel in Lemma 2.2.

Lemma 2.1. *Let $p_t(x, y) = p_t(a, x, y)$ be as in (2.16). There exists a random variable $\mathcal{X}$ such that for all $p \in [1, +\infty)$,*

$$(2.18) \quad \langle |\mathcal{X}|^p \rangle \leq C(p) < +\infty,$$

and

$$(2.19) \quad p_t(0, 0) \leq \frac{\mathcal{X}}{t^{d/2}}$$

or, equivalently

$$(2.20) \quad \sum_{x \in \mathbb{Z}^d} p_{t/2}(x, 0)^2 \leq \frac{\mathcal{X}}{t^{d/2}}.$$

Lemma 2.2. *For every $\alpha > \frac{d}{2} + 1$, there exists $C = C(\alpha, d) < +\infty$ such that for every $t \in \mathbb{R}^+$ it holds*

$$(2.21) \quad \omega_{\alpha-1}(t, x) p_t(x, 0) \leq C \frac{\sqrt{\bar{\mathcal{X}}(0) \bar{\mathcal{X}}(x)}}{t^{\frac{d}{2}}},$$

$$(2.22) \quad \sum_{b \in \mathbb{B}^d} \omega_\alpha^2(\underline{b}, t) |\sqrt{\bar{a}} \nabla p_t(b, 0)|^2 \leq C \frac{\mathcal{Y}_t}{(1+t)^{\frac{d}{2}+1}},$$

where $\bar{\mathcal{X}}(x) := \mathcal{X}(\tau_x a)$ is the stationary extension of the random variable $\mathcal{X}$ defined in Lemma 2.1 and

$$(2.23) \quad \mathcal{Y}_t := \mathcal{X}^{\frac{3}{2}} (1+t)^{-\frac{d}{2}} \sum_z \omega_{\frac{d}{2}+2}^{-2}(z, t) \sqrt{\bar{\mathcal{X}}(z)}.$$

Moreover, for every $p \in [1, +\infty)$ it holds

$$(2.24) \quad \langle |\sup_{t>0} \mathcal{Y}_t|^p \rangle^{\frac{1}{p}} \leq C(p, d) < +\infty.$$

The proof of Lemma 2.1 consists in showing that the environments we consider are “ w -moderate” in the sense defined in [10].

Lemma 2.3. *There exists a family of non-negative random variables $\{w(e)\}_{e \in \mathbb{B}}$ and a family of nearest-neighbor paths $\{\pi(e)\}_{e \in \mathbb{B}}$ such that the following properties hold:*

(i) *For every $e \in \mathbb{B}^d$ and $q \in [1, +\infty)$*

$$(2.25) \quad \langle |w(e)|^{-q} \rangle \leq C(d, q) < +\infty;$$

(ii) *Let $\xi : \Omega \rightarrow \mathbb{R}$; for every $e \in \mathbb{B}^d$, the path $\pi(e)$ connects the two endpoints of e and satisfies*

$$(2.26) \quad w(e)|D_e \xi|^2 \leq \sum_{b \in \pi(e)} a(b)|D_b \xi|^2.$$

Moreover, its length $|\pi(e)|$ is such that for every $q \in [1, +\infty)$,

$$(2.27) \quad \langle |\pi(e)|^q \rangle \leq C(d, q) < +\infty.$$

(iii) *Both $w(\cdot)$ and $\pi(\cdot)$ are stationary;*

Proof. In this proof the notation $\lesssim$ stands for $\leq C$ with $C = C(d, q)$. For every edge $e \in \mathbb{B}$, we define

$$(2.28) \quad w(e)^{-1} := \inf \left\{ \sum_{b \in \pi(e)} a^{-1}(b) : \pi(e) \text{ connects the two endpoints of } e \right\}.$$

Since a^{-1} is bounded from below, there exists a path that achieves the infimum above. We choose one according to a fixed, deterministic tie-breaking rule, and denote it by $\pi(e)$. With this definition of weights and paths, the point (iii) immediately follows by stationarity of a . We also have

$$\begin{aligned} |D_e \xi|^2 &= \left| \sum_{b \in \pi(e)} a(b)^{-\frac{1}{2}} a(b)^{\frac{1}{2}} D_b \xi \right|^2 \\ &\leq \left(\sum_{b \in \pi(e)} a(b)^{-1} \right) \left(\sum_{b \in \pi(e)} a(b) |D_b \xi|^2 \right) \stackrel{(2.28)}{=} w(e)^{-1} \left(\sum_{b \in \pi(e)} a(b) |D_b \xi|^2 \right), \end{aligned}$$

i.e. inequality (2.26). Moreover, since $a^{-1} \geq 1$, we have

$$|\pi(e)| \leq \sum_{b \in \pi(e)} a(b)^{-1},$$

and thanks to (2.28), the bound (2.27) is directly implied by (2.25). In order to show this last bound, we want to argue that for every $q \in [1, +\infty)$ and $x \gg 1$

$$(2.29) \quad \mathbb{P}(w^{-1}(e) > x) \lesssim x^{-q}.$$

We proceed in the following way: Thanks to assumption (1.3) and independence, it holds for $y \in \mathbb{R}$ to be fixed below

$$\mathbb{P}((\sup_i a(e_i))^{-1} \geq y) = \prod_{i=1}^d \mathbb{P}(a(e_i)^{-1} \geq y) \lesssim y^{-2qd},$$

and therefore there exists a (random) $i = i(y)$ such that

$$(2.30) \quad \mathbb{P}(a(e_{i(y)})^{-1} \geq y) \lesssim y^{-2q}.$$

The main idea is to explicitly construct a path $\tilde{\pi}(e)$, connecting the two endpoints of e for which we have some control on the quantity $\mathbb{P}(\sum_{b \in \tilde{\pi}(e)} a(b)^{-1} > x)$: From that, thanks to definition (2.28), we also obtain the same bound for (2.29). Without loss of generality, let us assume that $e = (z, z + e_1)$, for some $z \in \mathbb{Z}^d$. Therefore, if $i(y) = 1$ we just choose $\tilde{\pi}(e) = e$ and get by stationarity that for every $x > y$

$$(2.31) \quad \mathbb{P}(a(e)^{-1} > x) = \mathbb{P}(a(e)^{-1} > x) \stackrel{(2.30)}{\lesssim} y^{-2q},$$

i.e. the bound (2.25). If otherwise $i \neq 1$, then by stationarity and our assumption on the random variables $\{a(b)\}$ to be non-degenerate, we may fix a $0 < \varepsilon$ (independent on i and x) and consider

$$K := \inf\{k \geq 0 : a((z + k\bar{e}_i, z + k\bar{e}_i + \bar{e}_1) > \varepsilon\},$$

which satisfies

$$(2.32) \quad \mathbb{P}(K > k) \leq \exp(-ck),$$

for a positive constant $c = c(\varepsilon)$. Therefore, we estimate for any $x > \frac{2}{\varepsilon}$

$$(2.33) \quad \mathbb{P}(w^{-1}(e) \geq x) = \mathbb{P}(w^{-1}(e) \geq x, K > k) + \mathbb{P}(w^{-1}(e) \geq x, K \leq k).$$

We control the first term on the r.h.s of (2.33) by

$$(2.34) \quad \begin{aligned} \mathbb{P}(w^{-1}(e) \geq x, K \geq k) \\ = \mathbb{P}(w^{-1}(e) \geq x \mid K \geq k) \mathbb{P}(K \geq k) \stackrel{(2.32)}{\leq} \exp(-ck). \end{aligned}$$

For the second term the idea is to observe that, if $K \leq k$, then we might consider as path

$$\tilde{\pi}(e) := \{\tilde{b}_1, \dots, \tilde{b}_k, \tilde{e}, \tilde{b}_{k+1}, \dots, \tilde{b}_{2k}\}$$

the one starting from z , moving k steps in direction e_i , then moving in direction e_1 and finally going back with other k steps to $x + e_1$. Therefore,

$$\mathbb{P}(w^{-1}(e) \geq x, K < k) \leq \mathbb{P}\left(\sum_{b \in \tilde{\pi}(e)} a^{-1}(b) \geq x, K \leq k\right),$$

and since by construction

$$|\tilde{\pi}(e)| = 2k + 1 \quad \text{and} \quad \sum_{b \in \tilde{\pi}(e)} a(b)^{-1} \leq \varepsilon^{-1} \sum_{j=1}^{2k} a(\tilde{b}_j)^{-1},$$

we may control

$$\begin{aligned} \mathbb{P}(w^{-1}(e) \geq x, K < k) \\ \leq \mathbb{P}(\exists j \in \{1, \dots, 2k\} \text{ such that } a(\tilde{b}_j) \geq \frac{x}{4k}, K \leq k). \end{aligned}$$

Independence and then stationarity hence yield

$$\mathbb{P}(w^{-1}(e) \geq x, K < k) \leq 2k \mathbb{P}(a(e_i)^{-1} \geq \frac{x}{4k}).$$

Fixing now $k = x^\eta$ with $\eta < 1$, we get

$$\mathbb{P}(w^{-1}(e) \geq x, K < x^\eta) \leq x^\eta \mathbb{P}(a(e_i)^{-1} \geq \frac{x^{1-\eta}}{4}),$$

so that if we choose $y = \frac{x^{1-\eta}}{4} < x$ in (2.30), this turns into

$$\mathbb{P}(w^{-1}(e) \geq x, K < x^\eta) \lesssim x^{-q},$$

and (2.34) and (2.31) respectively into

$$\mathbb{P}(w^{-1}(e) \geq x, K > x^\eta) \leq \exp(-cx^\eta) \lesssim x^{-q},$$

$$\mathbb{P}(a(e)^{-1} > x) = \mathbb{P}(a(e)^{-1} > x) \lesssim x^{-2(1-\eta)q} \lesssim x^{-q}$$

By wrapping up the previous three inequalities we conclude (2.29) and hence (2.25). $\square$

Lemma 2.4. *For every $b \in \mathbb{B}$, let*

$$(2.35) \quad \pi^{-1}(b) := \{e \in \mathbb{B} : b \in \pi(e)\}.$$

Then, for every $p \in [1, +\infty)$

$$(2.36) \quad \langle |\pi^{-1}(b)|^p \rangle \leq C(d, p) < +\infty.$$

Proof. For a fixed edge $b \in \mathbb{B}^d$ and any $p \in [1, +\infty)$, let us consider

$$\mathbb{P}(|\pi^{-1}(b)| \geq k) = \mathbb{P}(|\pi^{-1}(b)| \geq k) = \mathbb{P}(\exists e_1, \dots, e_k : \forall i \quad b \in \pi(e_i)).$$

We observe that if there are $\sim k$ edges whose optimal path passes through b , then there must be an edge e with $|b - e| \geq k^{\frac{1}{d}}$. Therefore,

$$\mathbb{P}(|\pi^{-1}(b)| \geq k) \leq \mathbb{P}(\exists \tilde{e} \text{ with } |\tilde{e} - b| \geq k^{\frac{1}{d}} : \tilde{e} \in \pi(b)).$$

The path $\pi(\tilde{e})$ being connected, allows us to estimate

$$\begin{aligned} \mathbb{P}(|\pi^{-1}(b)| \geq k) &\leq \mathbb{P}(\exists \tilde{e} \text{ with } |\tilde{e} - b| \geq k^{\frac{1}{d}} : |\pi(\tilde{e})| > |b - \tilde{e}|) \\ &\leq \sum_{e: |b-e| > k^{\frac{1}{d}}} \mathbb{P}(|\pi(e)| > |b - e|) \simeq \sum_{n \sim k^{\frac{1}{d}}}^{+\infty} \sum_{|e-b|=n} \mathbb{P}(|\pi(e)| > n). \end{aligned}$$

Chebyshev's inequality yields for every $q \in [1, +\infty)$

$$\mathbb{P}(|\pi^{-1}(b)| \geq k) \leq \sum_{n \sim k^{\frac{1}{d}}}^{+\infty} \sum_{|e-b|=n} n^{-q} \mathbb{E}[|\pi(e)|^q] \stackrel{(2.27)-(iii)}{\leq} C(d, q) \sum_{n \sim k^{\frac{1}{d}}}^{+\infty} n^{d-1-q}.$$

We may now choose q big enough, e.g. $q = 2(p+1)d$, to conclude

$$\mathbb{P}(|\pi^{-1}(b)| \geq k) \leq C(d, 2(p+1)d) k^{-(2p+1)},$$

which implies inequality (2.36) for every $p \in [1, +\infty)$. $\square$

We now show the following general result on stationary random fields.

Lemma 2.5. *Let $\zeta : \Omega \times \mathbb{Z}^d \rightarrow \mathbb{R}$ be a stationary random field. Then for every $p \in [1, +\infty)$, edges $e_0, b_0 \in \mathbb{B}$ and $\delta > 0$ there exists a $C = C(d, p, \delta) < +\infty$ such that*

$$(2.37) \quad \langle |\sum_{b \in \pi(e_0)} \zeta(a, b)|^p \rangle^{\frac{1}{p}} \leq C \langle |\zeta(a, 0)|^{p(1+\delta)} \rangle^{\frac{1}{p(1+\delta)}},$$

$$(2.38) \quad \langle |\sum_{e \in \pi^{-1}(b_0)} \zeta(a, e)|^p \rangle^{\frac{1}{p}} \leq C \langle |\zeta(a, 0)|^{p(1+\delta)} \rangle^{\frac{1}{p(1+\delta)}},$$

whenever the r.h.s. of (2.37)-(2.38) is finite.

Proof. For the sake of simplicity, we skip the argument a in ζ and write $\lesssim$ instead of $\leq C$, with C depending on d, p and δ . We start with (2.38): Let us fix $p \in [1, +\infty)$ and $\delta > 0$. Since by (2.36) we have $\langle \cdot \rangle$ -almost surely that $|\pi^{-1}(z)| < +\infty$, we may

write

$$\begin{aligned}
& \langle | \sum_{e \in \pi^{-1}(b_0)} \zeta(\underline{e})|^p \rangle \\
&= \sum_{n=0}^{+\infty} \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n) \langle | \sum_{e \in \pi^{-1}(b_0)} \zeta(\underline{e})|^p | \max_{e \in \pi^{-1}(b_0)} |e - b_0| = n \rangle \\
&\lesssim \sum_{n=0}^{+\infty} \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n) \langle | \sum_{e \in B_{n+1}(\underline{b_0})} \zeta(\underline{e})|^p | \max_{e \in \pi^{-1}(b_0)} |e - b_0| = n \rangle,
\end{aligned}$$

and by Hölder inequality in e

$$\begin{aligned}
& \langle | \sum_{e \in \pi^{-1}(b_0)} \zeta(\underline{e})|^p \rangle \\
&\lesssim \sum_{n=0}^{+\infty} (n+1)^{d(p-1)} \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n) \sum_{e \in B_{n+1}(\underline{b_0})} \langle |\zeta(\underline{e})|^p | \max_{e \in \pi^{-1}(b_0)} |e - b_0| = n \rangle.
\end{aligned}$$

We now decompose the second term in the r.h.s. of the previous inequality as

$$\mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n) = \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n)^{\frac{\delta}{1+\delta}} \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n)^{\frac{1}{1+\delta}},$$

and thus rewrite

$$\begin{aligned}
& \langle | \sum_{e \in \pi^{-1}(b_0)} \zeta(\underline{e})|^p \rangle \\
&\lesssim \sum_{n=0}^{+\infty} (n+1)^{d(p-1)} \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n)^{\frac{\delta}{1+\delta}} \\
&\quad \times \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n)^{\frac{1}{1+\delta}} \sum_{e \in B_{n+1}(\underline{b_0})} \langle |\zeta(\underline{e})|^p | \max_{e \in \pi^{-1}(b_0)} |e - b_0| = n \rangle.
\end{aligned}$$

Therefore, an application of Hölder's inequality with exponents $(1 + 2\delta, \frac{1+2\delta}{2\delta})$ in n yields

$$\begin{aligned}
(2.39) \quad & \langle | \sum_{e \in \pi^{-1}(b_0)} \overline{\zeta(\underline{e})}|^p \rangle \lesssim \left(\sum_{n=0}^{+\infty} (n+1)^{d(p-1) \frac{1+2\delta}{2\delta}} \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n)^{\frac{1+2\delta}{2(1+\delta)}} \right)^{\frac{2\delta}{1+2\delta}} \\
& \times \left(\sum_{n=0}^{+\infty} \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n)^{1+\frac{\delta}{1+\delta}} \left(\sum_{e \in B_{n+1}(\underline{b_0})} \langle |\zeta(\underline{e})|^p | \max_{e \in \pi^{-1}(b_0)} |e - b_0| = n \rangle \right)^{1+2\delta} \right)^{\frac{1}{1+2\delta}}.
\end{aligned}$$

We now observe that the first term on the r.h.s. of (2.39) may be bounded by

$$(2.40) \quad \sum_{n=0}^{+\infty} (n+1)^{d(p-1) \frac{1+2\delta}{2\delta}} \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n)^{\frac{1+2\delta}{2(1+\delta)}} \lesssim 1.$$

This follows after noting that since

$$\mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n) \leq \mathbb{P}(\exists \tilde{e} \text{ with } |\tilde{e} - b| \geq n : |\pi(e)| > n),$$

by the same reasoning of Lemma 2.4 we have for every $M \geq 1$ that

$$(2.41) \quad \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n) \leq C(d, M) n^{-M},$$

and thus (2.40). We turn to the second term on the r.h.s. of (2.39) and claim that

$$(2.42) \quad \sum_{n=0}^{+\infty} \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n)^{1 + \frac{\delta}{1+\delta}} \left(\sum_{e \in B_{n+1}(\underline{b_0})} \langle |\zeta(\underline{e})|^p \mid \max_{e \in \pi^{-1}(b_0)} |e - b_0| = n \rangle \right)^{1+2\delta} \lesssim \langle |\zeta|^{p(1+2\delta)} \rangle.$$

Indeed, we write

$$\begin{aligned} & \sum_{n=0}^{+\infty} \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n)^{1 + \frac{\delta}{1+\delta}} \left(\sum_{e \in B_{n+1}(\underline{b_0})} \langle |\zeta(\underline{e})|^p \mid \max_{e \in \pi^{-1}(b_0)} |e - b_0| = n \rangle \right)^{1+2\delta} \\ &= \sum_n \sum_m I[m = n] \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = m)^{\frac{\delta}{1+\delta}} \\ & \quad \times \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n) \left(\sum_{e \in B_{m+1}(\underline{b_0})} \langle |\zeta(\underline{e})|^p \mid \max_{e \in \pi^{-1}(b_0)} |e - b_0| = n \rangle \right), \end{aligned}$$

and by Hölder's inequality first in the e -variable and then in $\langle \cdot \rangle$

$$\begin{aligned} & \sum_{n=0}^{+\infty} \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n)^{1 + \frac{\delta}{1+\delta}} \left(\sum_{e \in B_{n+1}(\underline{b_0})} \langle |\zeta(\underline{e})|^p \mid \max_{e \in \pi^{-1}(b_0)} |e - b_0| = n \rangle \right)^{1+2\delta} \\ & \leq \sum_n \sum_m I[m = n] m^{d \frac{1+2\delta}{2\delta}} \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = m)^{\frac{\delta}{1+\delta}} \\ & \quad \times \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n) \sum_{e \in B_{m+1}(\underline{b_0})} \langle |\zeta(\underline{e})|^{p(1+2\delta)} \mid \max_{e \in \pi^{-1}(b_0)} |e - b_0| = n \rangle \\ & \leq \sum_m m^{d \frac{1+2\delta}{2\delta}} \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = m)^{\frac{\delta}{1+\delta}} \\ & \quad \times \sum_{e \in B_{m+1}(\underline{b_0})} \sum_n \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n) \langle |\zeta(\underline{e})|^{p(1+2\delta)} \mid \max_{e \in \pi^{-1}(b_0)} |e - b_0| = n \rangle \\ & = \sum_m m^{d \frac{1+2\delta}{2\delta}} \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = m)^{\frac{\delta}{1+\delta}} \sum_{e \in B_{m+1}(\underline{b_0})} \langle |\zeta(\underline{e})|^{p(1+2\delta)} \rangle. \end{aligned}$$

Our assumption that ζ is stationary thus implies

$$\begin{aligned} & \sum_{n=0}^{+\infty} \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = n)^{1 + \frac{\delta}{1+\delta}} \left(\sum_{e \in B_{n+1}(\underline{b_0})} \langle |\zeta(\underline{e})|^p \mid \max_{e \in \pi^{-1}(b_0)} |e - b_0| = n \rangle \right)^{1+2\delta} \\ & \simeq \langle |\zeta(0)|^{p(1+2\delta)} \rangle \sum_m m^{d(1 + \frac{1+2\delta}{2\delta})} \mathbb{P}(\max_{e \in \pi^{-1}(b_0)} |e - b_0| = m)^{\frac{\delta}{1+\delta}}. \end{aligned}$$

Reasoning as for (2.40), we conclude inequality (2.42). Inserting estimates (2.40) and (2.42) in (2.39) yields inequality (2.38), after relabeling $\delta = 2\delta$.

We now prove (2.37) in an analogous way: Thanks to assumption (2.27), it holds the identity

$$\langle \mid \sum_{e \in \pi^{-1}(b_0)} \zeta(\underline{e})|^p \rangle = \sum_{n=1}^{+\infty} \mathbb{P}(|\pi(e_0)| = n) \langle \mid \sum_{e \in \pi^{-1}(b_0)} \zeta(\underline{e})|^p \mid |\pi(e_0)| = n \rangle.$$

We now reason exactly as in the argument for (2.38), this time relying directly on (2.27) and on Chebyshev's inequality to infer the analogous of (2.41), i.e. that for every $M \geq 1$

$$\mathbb{P}(|\pi(e_0)| = n) \leq C(d, M) n^{-M}.$$

□

Proof of Lemma 2.1. Thanks to Lemma 2.3, (ii) we may apply Theorem 3.2 of [10] and obtain that there exist $r > 0$ and $q > d$ such that

$$p_t(0, 0) \leq t^{-\frac{d}{2}} \left(\sup_r r^{-d} \sum_{|\bar{e}| \leq r} w^{-q}(e) \right)^r,$$

with $w(e)$ defined in Lemma 2.3. It thus only remains to prove that

$$\mathcal{X} := \left(\sup_r r^{-d} \sum_{|\bar{e}| \leq r} w^{-q}(e) \right)^r$$

satisfies (2.18). This follows from Lemma 2.3, (i) and the maximal function estimate ([10], Corollary A.2)

$$\langle |\sup_r r^{-d} \sum_{|\bar{e}| \leq r} w^{-q}(e)|^p \rangle^{\frac{1}{p}} \leq C(p) \langle |w^{-q}|^p \rangle^{\frac{1}{p}},$$

for every $p \in (1, +\infty]$. Estimate (2.20) follows from (2.19) thanks to the identity

$$p_t(0, 0) = \sum_z p_{\frac{t}{2}}^2(x, 0).$$

This, thanks to the symmetry of p_t , is in turn a particular case of

$$(2.43) \quad p_t(x, 0) = \sum_z p_s(x, z) p_{t-s}(z, 0),$$

with $x \in \mathbb{Z}^d$ and $s \in (0, t)$. To show (2.43) it suffices to observe that since for every $s > 0$ we have

$$\begin{cases} \partial_t p_{s+t} - \nabla \cdot \bar{a} \nabla p_{s+t} = 0 & t > 0 \\ p_s(x, 0) = p_s(x, 0), \end{cases}$$

then the representation formula implies the semigroup property

$$p_{s+t}(x, 0) = \sum_z p_t(x, z) p_s(z, 0),$$

which is equivalent to (2.43) if we relabel $t = t + s$. $\square$

Before proving Lemma 2.2, we state the following auxiliary result, whose proof we postpone to the appendix.

Lemma 2.6. *Let $\alpha > \frac{d}{2} + 1$ and let $\mathcal{Z}$ be a non-negative random variable such that for every $p \in [1, +\infty)$*

$$(2.44) \quad \langle |\mathcal{Z}|^p \rangle \leq C(p) < +\infty.$$

We then have

$$(2.45) \quad \langle |\sup_{t>0} \sum_{z \in \mathbb{Z}^d} \omega_\alpha^{-2}(z, t) \mathcal{Z}(\tau_z a)|^p \rangle \leq C(d, p, \alpha) < +\infty,$$

where the weight ω_α is defined as in (2.17).

Proof of Lemma 2.2. We prove Lemma 2.2 similarly to [5, Theorem 3]. We remark that, in contrast with Theorem 3, we do not need to prove an optimal decay in time for the weighted ℓ^{2p} -norm in space, and we replace inequality (173) by the stochastic bound (2.20) of Lemma 2.1.

We start by observing that by [10, Proposition 3.3 or 3.4], Lemma 2.1 implies for $\alpha > \frac{d}{2} + 1$

$$(2.46) \quad \sum_{x \in \mathbb{Z}^d} \omega_\alpha^2(x, t) p_t^2(x, 0) \lesssim \frac{\mathcal{X}}{t^{\frac{d}{2}}},$$

where here and in the rest of this proof $\lesssim$ stands for $\leq C(d, \alpha)$.

We start by upgrading inequality (2.46) to the bound (2.21): Since for every $t > 0$ and $s \in (0, t)$

$$(2.47) \quad \omega_\alpha(t, x) \lesssim \omega_\alpha(s, x - z) \omega_\alpha(t - s, z),$$

we may choose the value $s = \frac{t}{2}$ in (2.47) and in (2.43) of Lemma 2.1 and obtain

$$(2.48) \quad \begin{aligned} \omega_\alpha(t, x) p_t(x, 0) &\lesssim \sum_z \omega_\alpha\left(\frac{t}{2}, z - x\right) p_{\frac{t}{2}}(x, z) \omega_\alpha\left(\frac{t}{2}, z\right) p_{\frac{t}{2}}(z, 0) \\ &\lesssim \left(\sum_z \omega_\alpha^2\left(\frac{t}{2}, z - x\right) p_{\frac{t}{2}}^2(x, z) \right)^{\frac{1}{2}} \left(\sum_z \omega_\alpha^2\left(\frac{t}{2}, z\right) p_{\frac{t}{2}}^2(z, 0) \right)^{\frac{1}{2}}. \end{aligned}$$

By symmetry and stationarity of p_t , it holds

$$p_t(a, x, z) = p_t(a, z, x) = p_t(\tau_x a, z - x, 0),$$

so that inequality (2.48) turns into

$$\begin{aligned} \omega_\alpha(x, t) p_t(x, 0) &\lesssim \left(\sum_z \omega_\alpha^2\left(\frac{t}{2}, z - x\right) p_{\frac{t}{2}}^2(\tau_x a, z - x, 0) \right)^{\frac{1}{2}} \left(\sum_z \omega_\alpha^2\left(\frac{t}{2}, z\right) p_{\frac{t}{2}}^2(a, z, 0) \right)^{\frac{1}{2}} \\ &\stackrel{(2.20)}{\lesssim} \frac{\sqrt{\mathcal{X}(a) \mathcal{X}(\tau_x a)}}{t^{\frac{d}{2}}}. \end{aligned}$$

Recalling our definition of stationary extension of a random variable, the previous inequality yields (2.21).

In order to get also (2.22), we first claim that for every $T > 0$

$$(2.49) \quad \oint_T^{2T} \sum_b \omega_\alpha^2(\underline{b}, t) |\sqrt{\bar{a}(\underline{b})} \nabla p_t(\underline{b}, 0)|^2 \lesssim \frac{\mathcal{X}}{T^{\frac{d}{2}+1}}.$$

The identity

$$\frac{d}{dt} \left(\sum_x \frac{1}{2} p_t^2(x, 0) \right) \stackrel{(2.15)}{=} - \sum_b \nabla p_t(\underline{b}, 0) \cdot \bar{a}(\underline{b}) \nabla p_t(\underline{b}, 0)$$

implies by integrating

$$\oint_T^{2T} \sum_b \nabla p_t \cdot \bar{a} \nabla p_t(\underline{b}, 0) \lesssim T^{-1} \sum_x p_T^2(x, 0) \stackrel{(2.20)}{\lesssim} \frac{\mathcal{X}}{T^{\frac{d}{2}+1}}.$$

Therefore, as

$$\omega_\alpha^2(\underline{b}, t) \nabla p_t \cdot \bar{a} \nabla p_t(\underline{b}, 0) \lesssim t^{-\alpha} |\underline{b}|^{2\alpha} \nabla p_t \cdot \bar{a} \nabla p_t(\underline{b}, 0) + \nabla p_t \cdot \bar{a} \nabla p_t(\underline{b}, 0),$$

in order to show (2.49) it suffices to prove that

$$(2.50) \quad \oint_T^{2T} \sum_b |\underline{b}|^2 \nabla p_t(\underline{b}, 0) \cdot \bar{a}(\underline{b}) \nabla p_t(\underline{b}, 0) \lesssim \frac{\mathcal{X}}{T^{-\alpha+\frac{d}{2}+1}}.$$

We write

$$\begin{aligned} &\frac{d}{dt} \left(\sum_x |x|^{2\alpha} p_t^2(x, 0) \right) \stackrel{(2.15)}{=} - \sum_b \nabla (|\cdot|^{2\alpha} p_t(\cdot, 0))(\underline{b}) \cdot \bar{a}(\underline{b}) \nabla p_t(\underline{b}, 0) \\ &\lesssim - \sum_b |\underline{b}|^{2\alpha} \nabla p_t \cdot \bar{a} \nabla p_t(\underline{b}, 0) + \sum_b p_t(\bar{\underline{b}}, 0) (|\bar{\underline{b}}|^{2\alpha} - |\underline{b}|^{2\alpha}) |a(\underline{b}) \nabla p_t(\underline{b}, 0)| \\ &\lesssim - \sum_b |\underline{b}|^{2\alpha} \nabla p_t \cdot \bar{a} \nabla p_t(\underline{b}, 0) + \sum_b p_t(\bar{\underline{b}}, 0) |\bar{\underline{b}}|^{\alpha-1} |\underline{b}|^\alpha |a(\underline{b}) \nabla p_t(\underline{b}, 0)|, \end{aligned}$$

and thus by Hölder's inequality and Young's inequality we get

$$\frac{d}{dt} \left(\sum_x |x|^{2\alpha} p_t^2(x, 0) \right) \lesssim - \sum_b |\underline{b}|^{2\alpha} \nabla p_t \cdot a \nabla p_t(b, 0) + \sum_x |x|^{2(\alpha-1)} p_t^2(x, 0).$$

Integrating in $t \in (T, 2T)$ we obtain

$$\begin{aligned} \int_T^{2T} \sum_b |\underline{b}|^{2\alpha} \nabla p_t \cdot \bar{a} \nabla p_t(b, 0) \\ \lesssim T^{-1} \left(\sum_x |x|^{2\alpha} p_T^2(x, 0) \right) + \int_T^{2T} \sum_x |x|^{2(\alpha-1)} p_t^2(x, 0). \end{aligned}$$

Using (2.46) we conclude (2.50).

We finally prove (2.22): by (2.43) it holds for every $b \in \mathbb{B}$

$$\sqrt{\bar{a}} \nabla p_T(b, 0) = \int_{\frac{T}{3}}^{\frac{2}{3}T} \sum_z \sqrt{\bar{a}} \nabla p_t(b, z) p_{T-t}(z, 0) dt,$$

so that

$$\begin{aligned} \nabla p_T \cdot \bar{a} \nabla p_T(b, 0) &= |\sqrt{\bar{a}} \nabla p_T(b, 0)|^2 \\ &= \left| \int_{\frac{T}{3}}^{\frac{2}{3}T} \sum_z \sqrt{\bar{a}} \nabla p_t(b, z) p_{T-t}(z, 0) dt \right|^2 \\ (2.51) \quad &\leq \int_{\frac{T}{3}}^{\frac{2}{3}T} \sum_z p_{T-t}(z, 0) |\sqrt{\bar{a}} \nabla p_t(b, z)|^2 dt, \end{aligned}$$

where for the last line we appeal to Jensen's inequality in t and, thanks to $\sum_z p_s(z, 0) = 1$, also in z . By (2.47) we write

$$\begin{aligned} \sum_b \omega_\alpha^2(\underline{b}, T) \nabla p_T \cdot \bar{a} \nabla p_T(b, 0) \\ \stackrel{(2.47)}{\lesssim} \int_{\frac{T}{3}}^{\frac{2}{3}T} \sum_z \omega_\alpha^2(T-t, z) p_{T-t}(z, 0) \sum_b \omega_\alpha^2(t, \underline{b}-z) |\sqrt{\bar{a}} \nabla p_t(b, z)|^2 dt \\ \stackrel{(2.21), \beta > d}{\lesssim} \int_{\frac{T}{3}}^{\frac{2}{3}T} \sum_z \sqrt{\overline{\mathcal{X}(0)\mathcal{X}(z)}} (T-t)^{-\frac{d}{2}} \omega_\beta^{-2}(T-t, z) \sum_b \omega_\alpha^2(\underline{b}-z, t) |\sqrt{\bar{a}} \nabla p_t(b, z)|^2 dt \\ \lesssim T^{-\frac{d}{2}} \sum_z \sqrt{\overline{\mathcal{X}(0)\mathcal{X}(z)}} \omega_\beta^{-2}(T, z) \int_{\frac{T}{3}}^{\frac{2}{3}T} \sum_b \omega_\alpha^2(\underline{b}-z, t) |\sqrt{\bar{a}} \nabla p_t(b, z)|^2 dt \\ \stackrel{(2.49)}{\lesssim} \frac{\mathcal{X}}{T^{d+1}} \sum_z \sqrt{\overline{\mathcal{X}(0)\mathcal{X}(z)}} \omega_\beta^{-2}(T, z), \end{aligned}$$

which implies (2.22) if we choose $\beta = \frac{d}{2} + 2$.

To show inequality (2.24) we observe that thanks to (2.18) it is enough to prove that for every $p \in [1, +\infty)$

$$(2.52) \quad \langle |\sup_{t>0} (1+t)^{-\frac{d}{2}} \sum_z \omega_{\frac{d}{2}+2}^{-2}(z, t) \sqrt{\overline{\mathcal{X}(z)}}|^p \rangle \leq C(d, p) < +\infty.$$

This immediately follows from Lemma 2.6 since $\sqrt{\overline{\mathcal{X}(z)}}$ stands for $\sqrt{\overline{\mathcal{X}(a, z)}} = \sqrt{\overline{\mathcal{X}(\tau_z a)}}$ and we have chosen $\alpha = \frac{d}{2} + 2 > \frac{d}{2} + 1$. $\square$

3. PROOF OF THEOREM 1.1

Before giving the argument for Theorem 1.1 we introduce two technical results. The first is a generalization of Lemma 15 of [5].

Lemma 3.1. *Assume that*

$$(3.53) \quad 0 \leq a(t) \leq (1+t)^{-\gamma} + \int_0^t (1+t-s)^{-\gamma} b^\beta(s) ds,$$

$$(3.54) \quad 0 \leq b^p(t) \leq -\frac{d}{dt}[a^p(t)],$$

with $p \in [2, +\infty)$, $\gamma \in [1, +\infty)$, and $\beta \in (\frac{\gamma}{\gamma+1}, 1)$. Then there exists a constant $C = C(\lambda, \beta, p) < +\infty$ such that

$$(3.55) \quad a(t) \leq C(1+t)^{-\gamma}.$$

Proof. In this proof we use the notation $\lesssim$ for $\leq C(\lambda, \beta, p)$. We define

$$(3.56) \quad \Lambda(t) := \sup_{s \leq t} (1+s)^\gamma a(s).$$

By Hölder's inequality and (3.54), for any $0 \leq t_1 \leq t_2$ it follows

$$(3.57) \quad \int_{t_1}^{t_2} b^\beta(s) ds \leq \left(\int_{t_1}^{t_2} b^p(s) ds \right)^{\frac{\beta}{p}} (t_2 - t_1)^{1-\frac{\beta}{p}} \leq (t_2 - t_1)^{1-\frac{\beta}{p}} a^\beta(t_1).$$

Moreover, letting $N \in \mathbb{N}$ be such that $2^{N-1}t_1 \leq t_2 \leq 2^N t_1$, we get

$$\int_{t_1}^{t_2} b^\beta(s) ds \leq \sum_{n=1}^N \int_{2^{n-1}t_1}^{2^n t_1} b^\beta(s) ds \stackrel{(3.57)}{\leq} \sum_{n=1}^N t_1^{1-\frac{\beta}{p}} 2^{(n-1)(1-\frac{\beta}{p})} a^\beta(2^{n-1}t_1),$$

which by the definition of Λ in (3.56) implies

$$(3.58) \quad \begin{aligned} \int_{t_1}^{t_2} b^\beta(s) ds &\leq t_1^{1-\frac{\beta}{p}} \Lambda^\beta(t_2) \sum_{n=1}^N 2^{(n-1)(1-\frac{\beta}{p})} (2^{n-1}t_1)^{-\gamma\beta} \\ &\leq t_1^{1-\beta(\frac{1}{p}+\gamma)} \Lambda^\beta(t_2) \sum_{n=1}^N 2^{(n-1)(1-\beta(\frac{1}{p}+\gamma))} \lesssim \Lambda^\beta(t_2) t_1^{1-\beta(\frac{1}{p}+\gamma)}, \end{aligned}$$

since we assume $\beta(\frac{1}{p} + \gamma) > \gamma \geq 1$. Moreover,

$$\begin{aligned} a(t) &\stackrel{(3.54)}{\leq} \frac{1}{2t} \int_{\frac{t}{2}}^t a(r) dr \\ &\stackrel{(3.53)}{\leq} \frac{1}{2t} \int_{\frac{t}{2}}^t (1+r)^{-\gamma} dr + \frac{1}{2t} \int_{\frac{t}{2}}^t dr \int_0^r (1+r-s)^{-\gamma} b^\beta(s) ds \\ &\lesssim (1+t)^{-\gamma} + \frac{1}{2t} \int_{\frac{t}{2}}^t dr \int_0^r (1+r-s)^{-\gamma} b^\beta(s) ds. \end{aligned}$$

We let $\tau \in [0, \frac{t}{4}]$ to be chosen later, and write

$$\begin{aligned} a(t) &\lesssim (1+t)^{-\gamma} + \frac{1}{2t} \int_{\frac{t}{2}}^t dr \int_0^\tau (1+r-s)^{-\gamma} b^\beta(s) ds \\ &\quad + \frac{1}{2t} \int_{\frac{t}{2}}^t dr \int_\tau^{\frac{r}{2}} (1+r-s)^{-\gamma} b^\beta(s) ds \\ &\quad + \frac{1}{2t} \int_{\frac{t}{2}}^t dr \int_{\frac{r}{2}}^r (1+r-s)^{-\gamma} b^\beta(s) ds. \end{aligned}$$

We estimate each of the three last terms in turn:

$$\begin{aligned}
& \frac{1}{2t} \int_{\frac{t}{2}}^t dr \int_0^\tau (1+r-s)^{-\gamma} b^\beta(s) ds \lesssim (1+t)^{-\gamma} \int_0^\tau b^\beta(s) ds \\
& \stackrel{(3.57)}{\lesssim} (1+t)^{-\gamma} \tau^{1-\frac{\beta}{p}} a^\beta(0) \lesssim (1+t)^{-\gamma} \tau^{1-\frac{\beta}{p}}; \\
& \frac{1}{2t} \int_{\frac{t}{2}}^t dr \int_\tau^{\frac{r}{2}} (1+r-s)^{-\gamma} b^\beta(s) ds \\
& \lesssim (1+t)^{-\gamma} \int_\tau^{\frac{t}{2}} b^\beta(s) ds \stackrel{(3.58)}{\lesssim} (1+t)^{-\gamma} \frac{\Lambda^\beta(t)}{\tau^{\beta(\frac{1}{p}+\gamma)-1}}; \\
& \frac{1}{2t} \int_{\frac{t}{2}}^t dr \int_{\frac{r}{2}}^r (1+r-s)^{-\gamma} b^\beta(s) ds \\
& \lesssim \frac{1}{2t} \int_{\frac{t}{4}}^t b^\beta(s) ds \int_s^t (1+r-s)^{-\gamma} dr \\
& \stackrel{\gamma \geq 1}{\lesssim} \frac{1}{2t} \int_{\frac{t}{4}}^t b^\beta(s) ds \int_0^{t-s} (1+r)^{-1} dr \\
& \lesssim \left(\int_{\frac{t}{4}}^t b^\beta(s) ds \right) \left(\frac{1}{2t} \int_0^t (1+r)^{-1} dr \right) \\
& \lesssim \frac{\log(t+1)}{t} \int_{\frac{t}{4}}^t b^\beta(s) ds \stackrel{(3.58)}{\lesssim} \frac{\log(t+1)}{t} \frac{\Lambda^\beta(t)}{(1+t)^{\beta(\frac{1}{p}+\gamma)-1}} \\
& \lesssim \frac{\log(1+t)}{(t+1)^{\beta(\frac{1}{p}+\gamma)-\gamma}} \frac{\Lambda^\beta(t)}{(1+t)^\gamma}.
\end{aligned}$$

Everything together implies

$$(1+t)^\gamma a(t) \lesssim 1 + \tau^{1-\frac{\beta}{p}} + \left(\frac{1}{\tau^{\beta(\frac{1}{p}+\gamma)-1}} + \frac{\log(t+1)}{(t+1)^{\beta(\frac{1}{p}+\gamma)-\gamma}} \right) \Lambda^\beta(t),$$

where the implicit constant does not depend on our choice of $\tau \in [0, \frac{t}{4}]$. It follows from our assumption on β and γ that we can find $\tau_0 = \tau_0(\beta, \gamma, p) < \infty$ and $C(\beta, \gamma, p) < \infty$ such that for $\frac{t}{4} \geq \tau = \tau_0$,

$$(3.59) \quad (1+t)^\gamma a(t) \leq C \left(1 + \tau_0^{1-\frac{\beta}{p}} \right) + \frac{1}{2} \Lambda^\beta(t) \leq C \left(1 + \tau_0^{1-\frac{\beta}{p}} \right) + \frac{1}{2} \Lambda(t).$$

Hence, for such t , we have

$$\begin{aligned}
\Lambda(t) & \leq \sup_{0 \leq s \leq \tau_0} (1+s)^\gamma a(s) + \sup_{\tau_0 \leq s \leq t} (1+s)^\gamma a(s) \\
& \stackrel{(3.54), (3.59), 3.53}{\leq} (1+\tau_0)^\gamma + C \left(1 + \tau_0^{1-\frac{\beta}{p}} \right) + \frac{1}{2} \Lambda(t),
\end{aligned}$$

and this completes the proof. $\square$

Lemma 3.2. *Let u_t solve (1.6). Then, for every integer $p \geq 1$, we have*

$$(3.60) \quad \langle |Du_t \cdot a_t Du_t|^p \rangle \leq C \left(-\frac{d}{dt} \langle u^{2p}(t) \rangle \right),$$

where $C = C(d, p) < +\infty$.

Proof. We estimate

$$\frac{d}{dt}\langle u^{2p}(t) \rangle = -2p\langle Du^{2p-1} \cdot aDu \rangle$$

and we are done once that we prove

$$(3.61) \quad \langle Du^{2p-1} \cdot aDu \rangle \gtrsim \langle |Du \cdot aDu|^p \rangle,$$

where here and in the rest of this proof we write $\lesssim$ and $\gtrsim$ respectively for $\leq C$ and $\geq C$ with C depending on d and p . We note that the previous inequality can be rewritten as

$$\sum_{i=1}^d \langle D_i u^{2p-1} \cdot a_{ii} D_i u \rangle \gtrsim \langle \left| \sum_{i=1}^d D_i u \cdot a_{ii} D_i u \right|^p \rangle \gtrsim \sum_{i=1}^d \langle |D_i u \cdot a_{ii} D_i u|^p \rangle$$

so that we can reduce ourselves to prove for every fixed $i = 1, \dots, d$

$$D_i u^{2p-1} \cdot a_{ii} D_i u \gtrsim |D_i u \cdot a_{ii} D_i u|^p.$$

By our assumptions on the coefficients $0 \leq a_{ii} \leq 1$, we observe that if $a_{ii} = 0$, then the previous inequality is trivial. If otherwise $a_{ii} \neq 0$, then

$$(a^{2p-1} - 1)a_{ii}(a - 1) \stackrel{a_{ii} \leq 1}{\gtrsim} (a^{2p-1} - 1)a_{ii}^{2p}(a - 1),$$

and thus (3.61) is implied by

$$(3.62) \quad (a^{2p-1} - 1)(a - 1) \gtrsim (a - 1)^{2p},$$

for every $a \in \mathbb{R}$. We may now argue analogously to [5], Lemma 14, inequality (94) and show (3.62). $\square$

Proof of Theorem 1.1. Throughout this proof, the notation $\lesssim$ stands for $\leq C$ with the constant depending on d and p . We give ourselves an independent copy $(\tilde{a}(e))_{e \in \mathbb{B}}$ of the environment $(a(e))_{e \in \mathbb{B}}$, with the same law and defined on the same probability space. For each given $e \in \mathbb{B}$, we define the environment a^e by

$$(3.63) \quad a^e(b) := \begin{cases} a(b) & \text{if } b \neq e, \\ \tilde{a}(e) & \text{if } b = e. \end{cases}$$

In other words, the environment a^e is obtained from the environment a by resampling the conductance at the edge e .

By the independence assumption on the conductances $(a(e))$, every random variable $f \in L^2(\Omega)$ satisfies the spectral gap (or Efron-Stein) inequality

$$(3.64) \quad \langle (f - \langle f \rangle)^2 \rangle \leq \frac{1}{2} \sum_{e \in \mathbb{B}} \langle (\partial_e f)^2 \rangle,$$

with respect to the Glauber derivative

$$\partial_e f(a) := f(a^e) - f(a).$$

We start by observing that, analogously to Lemma 11 of [5], we can upgrade the spectral gap inequality (3.64) to

$$(3.65) \quad \langle |f - \langle f \rangle|^{2p} \rangle^{\frac{1}{p}} \lesssim \left\langle \left(\sum_e (\partial_e f)^2 \right)^p \right\rangle^{\frac{1}{p}}, \quad \text{for integer } p \in [1, +\infty).$$

Whenever no ambiguity occurs, we write $u_t(x) := \bar{u}_t(a, x)$ and/or skip the argument a in u and all the random variables involved. We now argue that,

appealing to (3.65), for u_t solution of (1.6) it holds for every integer $p \in [1, +\infty)$ that

$$(3.66) \quad \langle |u_t|^{2p} \rangle^{\frac{1}{p}} \lesssim \left\langle \left(\sum_e \left(\sum_z p_t(z, 0) \partial_e \bar{g}(z) \right)^2 \right)^p \right\rangle^{\frac{1}{p}} + \int_0^t \left\langle \left(\sum_b |\nabla p_{t-s}(b, 0)|^2 |\nabla \bar{u}_s(a^{e_i}, b)|^2 \right)^p \right\rangle^{\frac{1}{p}},$$

with $\bar{u}_t$, and $\bar{g}$ the stationary extensions solving (2.15).

To show the previous bound we take the Glauber derivative ∂_e in (2.15) and, thanks to the relation

$$[\partial_e, \nabla] := \partial_e \nabla - \nabla \partial_e = 0,$$

we obtain the parabolic boundary value problem

$$(3.67) \quad \begin{cases} \partial_t \partial_e \bar{u}_t + \nabla^* \bar{a} \nabla \partial_e \bar{u}_t = -\nabla^* h_t \\ \partial_e u_0 = \partial_e \bar{g}, \end{cases}$$

where $h_t = h_t(a, b, e)$ is defined as

$$(3.68) \quad h_t(a, b, e) := \partial_e \bar{a}(b) \nabla u_t(a^e, b) = \mathbf{1}_{b=e} \partial_e \bar{a}(e) \nabla u_t(a^e, e).$$

Using Duhamel's formula, equation (3.67) yields

$$\partial_e \bar{u}_t(x) = \sum_z p_t(z, x) \partial_e \bar{g}(z) + \int_0^t \left(\sum_b \nabla p_{t-s}(b, x) h_s(b, e) \right) ds,$$

which can be rewritten thanks to (3.68) as

$$(3.69) \quad \partial_e \bar{u}_t(x) = \sum_z p_t(z, x) \partial_e \bar{g}(z) + \int_0^t \left(\nabla p_{t-s}(e, x) \partial_e \bar{a}(e) \nabla u_s(a^e, e) \right) ds.$$

Furthermore, we note that for a general random variable $\zeta = \zeta(a)$ it holds

$$(3.70) \quad \begin{aligned} \partial_e \bar{\zeta}(a, x) &= \zeta((\tau_x a)^e) - \zeta(\tau_x a) \\ &= \zeta(\tau_x(a^{e-x})) - \zeta(\tau_x a) = \overline{\partial_{e-x} \zeta}(a, x), \end{aligned}$$

so that it follows for u_t that

$$\partial_e \bar{u}_t(a, 0) = \overline{\partial_e u_t}(a, 0).$$

Taking in the previous identity the $2p$ -th power and the average, stationarity implies that

$$(3.71) \quad \langle |\partial_e \bar{u}_t(0)|^{2p} \rangle = \langle |\partial_e u_t|^{2p} \rangle.$$

Since by (1.6) for both assumptions (a) and (b) on the initial data we have $\langle u_t \rangle = 0$, we may plug the identities (3.71) and (3.69) into the p-Spectral Gap (3.65) and get

$$(3.72) \quad \begin{aligned} \langle |u_t|^{2p} \rangle^{\frac{1}{p}} &\lesssim \left\langle \left(\sum_e (\partial_e \bar{u}_t(0))^2 \right)^p \right\rangle^{\frac{1}{p}} \\ &\lesssim \left\langle \left(\sum_e \left(\sum_z p_t(z, 0) \partial_e \bar{g}(z) \right)^2 \right)^p \right\rangle^{\frac{1}{p}} \\ &\quad + \left\langle \left(\sum_e \left(\int_0^t \nabla p_{t-s}(e, 0) \partial_e \bar{a}(e) \nabla u_s(a^e, e) ds \right)^2 \right)^p \right\rangle^{\frac{1}{p}}. \end{aligned}$$

We now apply the triangle inequality together with (3.68) on the second term on the r.h.s and estimate

$$\begin{aligned} & \left\langle \left(\sum_e \left(\int_0^t \left(\nabla p_{t-s}(e, 0) \partial_e \bar{a}(e) \nabla u_t(a^e, e) \right)^2 \right)^p \right)^{\frac{1}{p}} \right. \\ & \quad \left. \lesssim \int_0^t \left\langle \left(\sum_e |\nabla p_{t-s}(e, 0)|^2 |\nabla \bar{u}_s(a^e, e)|^2 \right)^p \right\rangle^{\frac{1}{p}}, \right. \end{aligned}$$

i.e. inequality (3.66).

We are now ready to prove part (a): We start focussing on the first term on the r.h.s. of (3.66) and claim that

$$(3.73) \quad \left\langle \left(\sum_e \left(\sum_z p_t(z, 0) \partial_e \bar{g}(z) \right)^2 \right)^p \right\rangle^{\frac{1}{p}} \lesssim t^{-\frac{d}{2}} N \|g\|_{L^\infty(\Omega)}^4.$$

We rewrite the l.h.s of (3.73) as

$$\begin{aligned} & \left\langle \left(\sum_{y, i=1, \dots, d} \left(\sum_z p_t(z, 0) \partial_{\{y, y+e_i\}} \bar{g}(z) \right)^2 \right)^p \right\rangle^{\frac{1}{p}} \\ & \quad \lesssim \sum_{i=1, \dots, d} \left\langle \left(\sum_y \left(\sum_z p_t(z, 0) \partial_{\{y, y+e_i\}} \bar{g}(z) \right)^2 \right)^p \right\rangle^{\frac{1}{p}}. \end{aligned}$$

For every fixed i , we estimate by the triangle inequality

$$\begin{aligned} & \left\langle \left(\sum_y \left(\sum_z p_t(z, 0) \partial_{\{y, y+e_i\}} \bar{g}(z) \right)^2 \right)^p \right\rangle^{\frac{1}{p}} \leq \sum_z \left\langle \left(\sum_y p_t^2(z, 0) |\partial_{\{y, y+e_i\}} \bar{g}(z)|^2 \right)^p \right\rangle^{\frac{1}{p}} \\ & \stackrel{(3.70)}{=} \sum_z \left\langle \left(\sum_y p_t^2(z, 0) |\overline{\partial_{\{y-z, y-z+e_i\}} g(z)}|^2 \right)^p \right\rangle^{\frac{1}{p}} \\ & \stackrel{(z=y-z)}{=} \sum_z \left\langle \left(\sum_y p_t^2(y-z, 0) |\overline{\partial_{\{z, z+e_i\}} g(y-z)}|^2 \right)^p \right\rangle^{\frac{1}{p}} \\ & \stackrel{y=y-z}{=} \sum_z \left\langle \left(\sum_y p_t^2(y, 0) |\overline{\partial_{\{z, z+e_i\}} g(y)}|^2 \right)^p \right\rangle^{\frac{1}{p}} \\ & \stackrel{(2.21)}{\lesssim} \sum_z \left\langle \left(\sum_y \sqrt{\mathcal{X}(0) \overline{\mathcal{X}(y)}} t^{-d} \omega_\alpha^{-2}(y, t) |\overline{\partial_{\{z, z+e_i\}} g(y)}|^2 \right)^p \right\rangle^{\frac{1}{p}}. \end{aligned} \quad (3.74)$$

We now use in the last term the triangle inequality in the inner sum and $\langle \cdot \rangle$ and infer that

$$\begin{aligned} & \left\langle \left(\sum_y \left(\sum_z p_t(z, 0) \partial_{\{y, y+e_i\}} \bar{g}(z) \right)^2 \right)^p \right\rangle^{\frac{1}{p}} \\ & \quad \lesssim t^{-d} \sum_z \sum_y \omega_\alpha^{-2}(y, t) \langle \sqrt{\mathcal{X}(0) \overline{\mathcal{X}(y)}}^{2p} |\overline{\partial_{\{z, z+e_i\}} g(y)}|^{2p} \rangle^{\frac{1}{p}}. \end{aligned}$$

After a repeated application of Hölder's inequality in $\langle \cdot \rangle$, stationarity and (2.18) yield

$$\begin{aligned} & \left\langle \left(\sum_y \left(\sum_z p_t(z, 0) \partial_{\{y, y+e_i\}} \bar{g}(z) \right)^2 \right)^p \right\rangle^{\frac{1}{p}} \\ & \lesssim t^{-d} \sum_z \|\partial_{\{z, z+e_i\}} g\|_{L^\infty(\Omega)}^4 \sum_y \omega_\alpha^{-2}(y, t) \\ & \stackrel{\alpha > \frac{d}{2}}{\lesssim} t^{-\frac{d}{2}} \sum_z \|\partial_{\{z, z+e_i\}} g\|_{L^\infty(\Omega)}^4. \end{aligned}$$

Appealing to our assumption on g to be in $L^\infty(\Omega)$ and depending on N edges, we get that

$$\left\langle \left(\sum_y \left(\sum_z p_t(z, 0) \partial_{\{y, y+e_i\}} \bar{g}(z) \right)^2 \right)^p \right\rangle^{\frac{1}{p}} \lesssim N \|g\|_{L^\infty(\Omega)}^2 t^{-\frac{d}{2}}.$$

Summing over $i = 1, \dots, d$ yields (3.73).

We now turn to the second term of the r.h.s. of (3.66) to argue that for every $\delta > 0$ it holds

$$\begin{aligned} & \int_0^t \left\langle \left(\sum_e |\nabla p_{t-s}(e, 0)|^2 |\nabla \bar{u}_s(a^e, e)|^2 \right)^p \right\rangle^{\frac{1}{p}} ds \\ (3.75) \quad & \lesssim C(\delta) \int_0^t (t-s)^{-(\frac{d}{2}+1)(1-\frac{1}{p})+\frac{d}{p}} \langle |Du_t \cdot a Du_t|^{p(1+\delta)} \rangle^{\frac{1}{p(1+\delta)}} ds. \end{aligned}$$

By (2.26) of Lemma 2.3, we write

$$\begin{aligned} & \int_0^t \left\langle \left(\sum_e |\nabla p_{t-s}(e, 0)|^2 |\nabla \bar{u}_s(a^e, e)|^2 \right)^p \right\rangle^{\frac{1}{p}} ds \\ & \lesssim \int_0^t \left\langle \left(\sum_e w(e)^{-1} \sum_{b \in \pi(e)} |\sqrt{\bar{a}(b)} \nabla p_{t-s}(b, 0)|^2 w'(e)^{-1} \sum_{b' \in \pi'(e)} |\sqrt{\bar{a}^e(b')} \nabla \bar{u}_s(a^e, b')|^2 \right)^p \right\rangle^{\frac{1}{p}} ds \\ & \simeq \int_0^t \left\langle \left(\sum_b |\sqrt{\bar{a}(b)} \nabla p_{t-s}(b, 0)|^2 \sum_{e \in \pi^{-1}(b)} w(e)^{-1} w'(e)^{-1} \sum_{b' \in \pi'(e)} |\sqrt{\bar{a}^e(b')} \nabla \bar{u}_s(a^e, b')|^2 \right)^p \right\rangle^{\frac{1}{p}} ds, \end{aligned}$$

where $\pi'(e)$ and $w'(e)$ are as π and w introduced in Lemma 2.3 and related to the environment given by a^e . After smuggling the weight $\omega_\alpha^2(\underline{b}, t-s)$ in the sum over b , we use Hölder's inequality with exponents q and p to estimate

$$\begin{aligned} & \int_0^t \left\langle \left(\sum_e |\nabla p_{t-s}(e, 0)|^2 |\nabla \bar{u}_s(a^e, e)|^2 \right)^p \right\rangle^{\frac{1}{p}} ds \\ & \lesssim \int_0^t \left\langle \left(\sum_b \omega_\alpha^{2q}(\underline{b}, t-s) |\sqrt{\bar{a}(b)} \nabla p_{t-s}(b, 0)|^{2q} \right)^{p-1} \right. \\ & \quad \times \left. \left(\sum_b \omega_\alpha^{-2p}(\underline{b}, t-s) \left(\sum_{e \in \pi^{-1}(b)} w(e)^{-1} w'(e)^{-1} \sum_{b' \in \pi'(e)} |\sqrt{\bar{a}^e(b')} \nabla \bar{u}_s(a^e, b')|^2 \right)^p \right) \right\rangle^{\frac{1}{p}} ds. \end{aligned}$$

We now appeal to Lemma 2.2 and the embedding $\ell^2 \subseteq \ell^r$ for $r \geq 2$ to infer from the previous inequality that

$$\begin{aligned}
& \int_0^t \left\langle \left(\sum_e |\nabla p_{t-s}(e, 0)|^2 |\nabla \bar{u}_s(a^e, e)|^2 \right)^p \right\rangle^{\frac{1}{p}} ds \\
& \lesssim \int_0^t (t-s)^{-(\frac{d}{2}+1)(1-\frac{1}{p})} \left(\sum_b \omega_\alpha^{-2p}(\underline{b}, t-s) \right. \\
(3.76) \quad & \left. \times \langle \mathcal{Y}_{t-s}^{p-1} \left(\sum_{e \in \pi^{-1}(b)} w(e)^{-1} w'(e)^{-1} \sum_{b' \in \pi'(z)} |\sqrt{\bar{a}^e(b')} \nabla \bar{u}_s(a^e, b')|^2 \right)^p \rangle \right)^{\frac{1}{p}} ds.
\end{aligned}$$

We claim that

$$\begin{aligned}
& \langle \mathcal{Y}_{t-s}^{p-1} \left(\sum_{e \in \pi^{-1}(b)} w(e)^{-1} w'(e)^{-1} \sum_{b' \in \pi'(e)} |\sqrt{\bar{a}^e(b')} \nabla \bar{u}_s(a^e, b')|^2 \right)^p \rangle \\
(3.77) \quad & \lesssim C(\delta) \langle |Du_s \cdot a Du_s|^{p(1+\delta)} \rangle^{\frac{1}{1+\delta}}.
\end{aligned}$$

We note that, once that we have (3.77), by inserting it in (3.76) we get

$$\begin{aligned}
& \int_0^t \left\langle \left(\sum_e |\nabla p_{t-s}(e, 0)|^2 |\nabla \bar{u}_s(a^e, e)|^2 \right)^p \right\rangle^{\frac{1}{p}} ds \\
& \lesssim C(\delta) \int_0^t (t-s)^{-(\frac{d}{2}+1)(1-\frac{1}{p})} \left(\sum_b \omega_\alpha^{-2p}(\underline{b}, t-s) \langle |Du_s \cdot a Du_s|^{p(1+\delta)} \rangle^{\frac{1}{1+\delta}} \right)^{\frac{1}{p}} ds \\
& \stackrel{\alpha > \frac{d}{2}}{\lesssim} C(\delta) \int_0^t (t-s)^{-(\frac{d}{2}+1)(1-\frac{1}{p}) + \frac{d}{p}} \langle |Du_s \cdot a Du_s|^{p(1+\delta)} \rangle^{\frac{1}{p(1+\delta)}} ds
\end{aligned}$$

and thus (3.75).

We now prove (3.77): It holds

$$\begin{aligned}
& \langle \mathcal{Y}_{t-s}^{p-1} \left(\sum_{e \in \pi^{-1}(b)} w(e)^{-1} w'(e)^{-1} \sum_{b' \in \pi'(e)} |\sqrt{\bar{a}^e(b')} \nabla \bar{u}_s(a^e, b')|^2 \right)^p \rangle \\
& \leq \langle \mathcal{Y}_{t-s}^{p-1} |\pi^{-1}(b)|^{p-1} \sum_{e \in \pi^{-1}(b)} w(e)^{-p} w'(e)^{-p} \left(\sum_{b' \in \pi'(e)} |\sqrt{\bar{a}^e(b')} \nabla \bar{u}_s(a^e, b')|^2 \right)^p \rangle \\
& = \langle \sum_{e \in \pi^{-1}(b)} \mathcal{Y}_{t-s}^{p-1} |\pi^{-1}(b)|^{p-1} w(e)^{-p} w'(e)^{-p} \left(\sum_{b' \in \pi'(e)} |\sqrt{\bar{a}^e(b')} \nabla \bar{u}_s(a^e, b')|^2 \right)^p \rangle.
\end{aligned}$$

Hölder's inequality with exponents $(1+\delta, \frac{1+\delta}{\delta})$ first in e and then in $\langle \cdot \rangle$ yields

$$\begin{aligned}
& \langle \mathcal{Y}_{t-s}^{p-1} \left(\sum_{e \in \pi^{-1}(b)} w(e)^{-1} w'(e)^{-1} \sum_{b' \in \pi'(e)} |\sqrt{\bar{a}^e(b')} \nabla \bar{u}_s(a^e, b')|^2 \right)^p \rangle \\
& \leq \langle \sum_{e \in \pi^{-1}(b)} \mathcal{Y}_{t-s}^{(p-1)\frac{1+\delta}{\delta}} |\pi^{-1}(b)|^{(p-1)\frac{1+\delta}{\delta}} w(e)^{-p\frac{1+\delta}{\delta}} w'(e)^{-p\frac{1+\delta}{\delta}} \rangle^{\frac{\delta}{1+\delta}} \\
(3.78) \quad & \times \langle \sum_{e \in \pi^{-1}(b)} \left(\sum_{b' \in \pi'(e)} |\sqrt{\bar{a}^e(b')} \nabla \bar{u}_s(a^e, b')|^2 \right)^{p(1+\delta)} \rangle^{\frac{1}{1+\delta}}
\end{aligned}$$

As the terms $\sum_{b' \in \pi'(e)} |\sqrt{\bar{a}^e(b')} \nabla \bar{u}_s(a^e, b')|^2$ and $\sqrt{\bar{a}^e(b')} \nabla \bar{u}_s(a^e, b')$ are stationary respectively by Lemma 2.3 and by definition of stationary extension, we appeal to

Lemma 2.5 and obtain that for every $\delta > 0$

$$\begin{aligned}
& \left\langle \sum_{e \in \pi^{-1}(b)} \left(\sum_{b' \in \pi'(e)} |\sqrt{\bar{a}^e(b')} \nabla \bar{u}_s(a^e, b')|^2 \right)^{p(1+\delta)} \right\rangle^{\frac{1}{1+\delta}} \\
& \stackrel{(2.38)}{\lesssim} C(\delta) \sum_{i=1}^d \left\langle \left(\sum_{b' \in \pi'(e_i)} |\sqrt{\bar{a}^{e_i}(b')} \nabla \bar{u}_s(a^{e_i}, b')|^2 \right)^{p(1+\delta)^2} \right\rangle^{\frac{1}{(1+\delta)^2}} \\
& \stackrel{(2.37)}{\lesssim} C(\delta) \sum_{i=1}^d \left\langle |\sqrt{\bar{a}^{e_i}} Du_s(a^{e_i})|^{2p(1+\delta)^3} \right\rangle^{\frac{1}{(1+\delta)^3}}.
\end{aligned}$$

Since a^e and a have, by definition, the same law we conclude that

$$\begin{aligned}
& \left\langle \sum_{e \in \pi^{-1}(b)} \left(\sum_{b' \in \pi'(e)} |\sqrt{\bar{a}^e(b')} \nabla \bar{u}_s(a^e, b')|^2 \right)^{p(1+\delta)} \right\rangle^{\frac{1}{1+\delta}} \\
(3.79) \quad & \lesssim C(\delta) \left\langle |\sqrt{\bar{a}} Du_s(a)|^{2p(1+\delta)^3} \right\rangle^{\frac{1}{(1+\delta)^3}}.
\end{aligned}$$

We now turn to the other averaged term in (3.78): Reasoning as above, thanks to (iii) of Lemma 2.3 and (2.23) of Lemma 2.2, we may apply (2.37) of Lemma 2.5 and get that for every $\delta > 0$ the last term on the r.h.s. of (3.78) satisfies

$$\begin{aligned}
& \left\langle \sum_{e \in \pi^{-1}(b)} \mathcal{Y}_{t-s}^{(p-1)\frac{1+\delta}{\delta}} |\pi^{-1}(b)|^{(p-1)\frac{1+\delta}{\delta}} w(e)^{-p\frac{1+\delta}{\delta}} w'(e)^{-p\frac{1+\delta}{\delta}} \right\rangle \\
& \lesssim \sum_{i=1}^d \left\langle \left(\mathcal{Y}_{t-s}^{(p-1)\frac{1+\delta}{\delta}} |\pi^{-1}(e_i)|^{(p-1)\frac{1+\delta}{\delta}} w(e_i)^{-p\frac{1+\delta}{\delta}} w'(e_i)^{-p\frac{1+\delta}{\delta}} \right)^{1+\delta} \right\rangle^{\frac{1}{1+\delta}} \\
& \stackrel{(2.25)-(2.27)-(2.24)}{\lesssim} C(\delta).
\end{aligned}$$

Inequalities (3.79) and (3.80) in (3.78) yield the bound (3.77), once that we relabel $\delta \simeq \delta^3 + 2\delta^2 + 2\delta$.

We are now ready to conclude the proof of Theorem 1.1, part (a): By (3.73) and (3.75), we get from (3.66)

$$\begin{aligned}
& \langle |u_t|^{2p} \rangle^{\frac{1}{p}} \lesssim C(\delta) N \|g\|_{L^\infty(\Omega)}^2 \left(t^{-\frac{d}{2}} \right. \\
& \quad \left. + \int_0^t (t-s)^{-(\frac{d}{2}+1)(1-\frac{1}{p})+\frac{d}{p}} \langle |Du_s \cdot a Du_s|^{p(1+\delta)} \rangle^{\frac{1}{p(1+\delta)}} ds \right),
\end{aligned}$$

and, by our assumption $f \in L^\infty(\Omega)$, that

$$\langle |u_t|^{2p} \rangle^{\frac{1}{p}} \lesssim C(\delta) N \|g\|_{L^\infty(\Omega)}^2 \left(t^{-\frac{d}{2}} + \int_0^t (t-s)^{-(\frac{d}{2}+1)(1-\frac{1}{p})+\frac{d}{p}} \langle |Du_t \cdot a Du_t|^p \rangle^{\frac{1}{p(1+\delta)}} ds \right).$$

If we now choose $p \in [1, +\infty)$ such that $(\frac{d}{2}+1)(1-\frac{1}{p})-\frac{d}{p} > \frac{d}{2}$ and $\delta > 0$, it follows that

$$\langle |u_t|^{2p} \rangle^{\frac{1}{p}} \lesssim C(\delta) N \|g\|_{L^\infty(\Omega)}^2 \left(t^{-\frac{d}{2}} + \int_0^t (t-s)^{-\frac{d}{2}} \langle |Du_t \cdot a Du_t|^p \rangle^{\frac{1}{p(1+\delta)}} ds \right).$$

and, in view of Lemma 3.2 and Lemma 3.1 with $\gamma := \frac{d}{2}$ any $\beta := \frac{1}{1+\delta} > \frac{\gamma}{\gamma+\frac{1}{p}}$, also (1.7) for p large enough. The embedding

$$(3.81) \quad \langle |u_t|^q \rangle^{\frac{1}{q}} \leq C(p, q) \langle |u_t|^p \rangle^{\frac{1}{p}}$$

allows to conclude the proof of Theorem 1.7, part (a).

To prove also Theorem 1.1, part (b) we observe that thanks to the assumption $g = D^*f$ (and thus $\bar{g} = \nabla^* \bar{f}$), after an integration by parts, we may reformulate the estimate (3.66) as

$$(3.82) \quad \langle |u_t|^{2p} \rangle^{\frac{1}{p}} \lesssim \left\langle \left(\sum_e \left(\sum_b \nabla p_t(b, 0) \partial_e \bar{f}(b) \right)^2 \right)^p \right\rangle^{\frac{1}{p}} + \int_0^t \left\langle \left(\sum_e |\nabla p_{t-s}(e, 0)|^2 |\nabla \bar{u}_s(a^e, e)|^2 \right)^p \right\rangle^{\frac{1}{p}}.$$

We control the second term on the r.h.s. again by (3.75). For the first term, we now argue that

$$(3.83) \quad \left\langle \left(\sum_e \left(\sum_b \nabla p_t(b, 0) \partial_e \bar{f}(b) \right)^2 \right)^p \right\rangle^{\frac{1}{p}} \lesssim N \|f\|_{L^\infty(\Omega)}^2 t^{-(\frac{d}{2}+1)(1-\frac{1}{p})+\frac{d}{p}},$$

with N the number of sites on which f depends.

Before proving (3.83), we show how to conclude the argument: Inserting inequalities (3.83) and (3.73) in (3.82), we get

$$\langle |u_t|^{2p} \rangle^{\frac{1}{p}} \lesssim C(\delta) N \|f\|_{L^\infty(\Omega)}^2 \left(t^{-(\frac{d}{2}+1)(1-\frac{1}{p})+\frac{d}{p}} + \int_0^t (t-s)^{-(\frac{d}{2}+1)(1-\frac{1}{p})+\frac{d}{p}} \langle |Du_t \cdot a Du_t|^p \rangle^{\frac{1}{p(1+\delta)}} ds \right),$$

so that by Lemma 3.2 and Lemma 3.1 with $\gamma := (\frac{d}{2} + 1)(1 - \frac{1}{p}) - \frac{d}{p}$ and any $\beta := \frac{1}{1+\delta} > \frac{\gamma}{\gamma+\frac{1}{p}}$, we obtain

$$\langle |u_t|^{2p} \rangle^{\frac{1}{p}} \lesssim N \|f\|_{L^\infty(\Omega)}^2 t^{-(\frac{d}{2}+1)(1-\frac{1}{p})+\frac{d}{p}} \lesssim N \|f\|_{L^\infty(\Omega)}^2 t^{-(\frac{d}{2}+1)+\frac{d}{2p}}.$$

Therefore, for every $q \in [1, +\infty)$ fixed, let us consider $\varepsilon > 0$; we choose $q \leq p < +\infty$ such that $\frac{d}{2p} \leq \varepsilon$ and use the previous estimate together with the embedding (3.81) to infer

$$\langle |u_t|^{2q} \rangle^{\frac{1}{q}} \lesssim C(\varepsilon) N \|f\|_{L^\infty(\Omega)}^2 t^{-(\frac{d}{2}+1)+\frac{d}{2p}} \lesssim C(\varepsilon) N \|f\|_{L^\infty(\Omega)}^2 t^{-(\frac{d}{2}+1)+\varepsilon},$$

i.e. bound (1.8).

It thus only remains to show (3.83): As in (3.74), we bound the l.h.s of (3.83) by

$$(3.84) \quad \left\langle \left(\sum_e \left(\sum_b \nabla p_t(b, 0) \partial_e \bar{f}(z) \right)^2 \right)^p \right\rangle^{\frac{1}{p}} \leq \sum_{i=1, \dots, d} \sum_z \left\langle \left(\sum_b |\nabla p_t(b, 0)|^2 |\bar{\partial}_{\{z, z+e_i\}} \bar{f}(b)|^2 \right)^p \right\rangle^{\frac{1}{p}}.$$

For any fixed $i = 1, \dots, d$, inequality (2.26) of Lemma 2.3 yields

$$\begin{aligned} & \sum_z \left\langle \left(\sum_b |\nabla p_t(b, 0)|^2 |\bar{\partial}_{\{z, z+e_i\}} \bar{f}(b)|^2 \right)^p \right\rangle^{\frac{1}{p}} \\ & \leq \sum_z \left\langle \left(\sum_b w(b)^{-1} \sum_{b' \in \pi(b)} |\sqrt{\bar{a}(b')} \nabla p_t(b', 0)|^2 |\bar{\partial}_{\{z, z+e_i\}} \bar{f}(b)|^2 \right)^p \right\rangle^{\frac{1}{p}} \\ & = \sum_z \left\langle \left(\sum_{b'} |\sqrt{\bar{a}(b')} \nabla p_t(b', 0)|^2 \sum_{b \in \pi^{-1}(b')} w(b)^{-2} |\bar{\partial}_{\{z, z+e_i\}} \bar{f}(b)|^2 \right)^p \right\rangle^{\frac{1}{p}} \end{aligned}$$

After smuggling the weight $\omega_\alpha^2(\underline{b}', t)$ into the sum over b' , Hölder's inequality with exponents p and q implies

$$\begin{aligned} \sum_z \left\langle \left(\sum_b |\nabla p_t(b, 0)|^2 |\overline{\partial_{\{z, z+e_i\}} f(b)}|^2 \right)^p \right\rangle^{\frac{1}{p}} \\ \lesssim \sum_z \left\langle \left(\sum_{b'} \omega_\alpha^{2q}(\underline{b}', t) |\sqrt{a(b')} \nabla p_t(b', 0)|^{2q} \right)^{p-1} \right. \\ \left. \times \left(\sum_{b'} \omega_\alpha^{-2p}(\underline{b}', t) \left(\sum_{b \in \pi^{-1}(b')} w(b)^{-1} |\overline{\partial_{\{z, z+e_i\}} f(b)}|^2 \right)^p \right) \right\rangle^{\frac{1}{p}}, \end{aligned}$$

and thus by Lemma 2.2,

$$\begin{aligned} \sum_z \left\langle \left(\sum_b |\nabla p_t(b, 0)|^2 |\overline{\partial_{\{z, z+e_i\}} f(b)}|^2 \right)^p \right\rangle^{\frac{1}{p}} \\ (3.85) \quad \lesssim t^{-(\frac{d}{2}+1)(1-\frac{1}{p})} \sum_z \left\langle \left(\sum_{b'} \omega_\alpha^{-2p}(\underline{b}', t) \left(\sum_{b \in \pi^{-1}(b')} w(b)^{-1} |\overline{\partial_{\{z, z+e_i\}} f(b)}|^2 \right)^p \right) \right\rangle^{\frac{1}{p}}. \end{aligned}$$

We now show that

$$(3.86) \quad \left\langle \left(\sum_{b \in \pi^{-1}(b')} w(b)^{-1} |\overline{\partial_{\{z, z+e_i\}} f(b)}|^2 \right)^p \right\rangle \lesssim \|\partial_{\{z, z+e_i\}} f\|_{L^\infty(\Omega)}^{2p}.$$

Once proven the previous inequality, estimate (3.85) turns into

$$\begin{aligned} \sum_z \left\langle \left(\sum_b |\nabla p_t(b, 0)|^2 |\overline{\partial_{\{z, z+e_i\}} f(b)}|^2 \right)^p \right\rangle^{\frac{1}{p}} \\ \lesssim t^{-(\frac{d}{2}+1)(1-\frac{1}{p})} \sum_z \left\langle \left(\sum_{b'} \omega_\alpha^{-2p}(\underline{b}', t) \right)^{\frac{1}{p}} \|\partial_{\{z, z+e_i\}} f\|_{L^\infty(\Omega)}^2 \right\rangle^{\frac{1}{p}} \\ \lesssim t^{-(\frac{d}{2}+1)(1-\frac{1}{p}) + \frac{d}{p}} \sum_z \|\partial_{\{z, z+e_i\}} f\|_{L^\infty(\Omega)}^2 \\ \lesssim N t^{-(\frac{d}{2}+1)(1-\frac{1}{p}) + \frac{d}{p}} \|f\|_{L^\infty(\Omega)}^2, \end{aligned}$$

where for the last inequality we observe that $\partial_{z, z+e_i} f = 0$ if $(z, z+e_i) \notin \text{supp}(f)$. The bound (3.83) follows by (3.84).

To obtain (3.86), we reason similarly to (3.79) and (3.80): We write

$$\begin{aligned} \left\langle \left(\sum_{b \in \pi^{-1}(b')} w(b)^{-1} |\overline{\partial_{\{z, z+e_i\}} f(b)}|^2 \right)^p \right\rangle \\ \stackrel{(2.37)}{\lesssim} \sum_{j=1}^d C(\delta) \langle w(e_j)^{-p(1+\delta)} |\overline{\partial_{\{z, z+e_i\}} f(e_j)}|^{2p(1+\delta)} \rangle^{\frac{1}{1+\delta}} \\ \stackrel{\delta=1}{\lesssim} \sum_{j=1}^d \langle w(e_j)^{-2p} |\overline{\partial_{\{z, z+e_i\}} f(e_j)}|^{4p} \rangle^{\frac{1}{2}}. \end{aligned}$$

Appealing to the assumption $f \in L^\infty(\Omega)$ and to the bound (2.25) of Lemma 2.3, we infer (3.86). $\square$

4. APPENDIX

Proof of Lemma 2.6. We start observing that it is enough to show an analogue of the maximal function estimate ([1], Theorem 3.2), namely that

$$(4.87) \quad \mathbb{P}(|\sup_{t>0} t^{-\frac{d}{2}} \sum_z \omega_\alpha^{-2}(z, t) \mathcal{Z}(\tau_z a)| > \lambda) \lesssim \frac{1}{\lambda} \langle |\mathcal{Z}| \rangle.$$

This indeed, combined with the Marcinkiewicz Interpolation Theorem and the fact that the map

$$\mathcal{Z} \rightarrow \sup_{t>0} t^{-\frac{d}{2}} \sum_z \omega_\alpha^{-2}(z, t) \mathcal{Z}(\tau_z a)$$

is bounded from L^∞ to L^∞ , yields for every $p \in (1, +\infty]$

$$\langle |\sup_{t>0} t^{-\frac{d}{2}} \sum_z \omega_\alpha^{-2}(z, t) \mathcal{Z}(\tau_z a)|^p \rangle \lesssim \langle |\mathcal{Z}|^p \rangle$$

and thus also (2.45) by assumption (2.44).

We now give the argument for (4.87): We observe that it holds

$$\begin{aligned} & \mathbb{P}(\sup_{t>0} t^{-\frac{d}{2}} \sum_z \omega_\alpha^{-2}(z, t) \mathcal{Z}(\tau_z a) > \lambda) \\ &= \mathbb{P}(\sup_{t>0} t^{-\frac{d}{2}} \sum_{n=1}^{\infty} \sum_{(n-1)t^{\frac{1}{2}} \leq |z| < nt^{\frac{1}{2}}} \omega_\alpha^{-2}(z, t) \mathcal{Z}(\tau_z a) > \lambda) \\ &= \mathbb{P}(\exists t \in (0, +\infty) : t^{-\frac{d}{2}} \sum_{n=1}^{\infty} \sum_{(n-1)t^{\frac{1}{2}} \leq |z| < nt^{\frac{1}{2}}} \omega_\alpha^{-2}(z, t) \mathcal{Z}(\tau_z a) > \lambda) \\ &= \mathbb{P}(\exists t \in (0, +\infty), n \in \mathbb{N} : t^{-\frac{d}{2}} \sum_{(n-1)t^{\frac{1}{2}} \leq |z| < nt^{\frac{1}{2}}} \omega_\alpha^{-2}(z, t) \mathcal{Z}(\tau_z a) \gtrsim \frac{\lambda}{n^2}). \end{aligned}$$

Since $\mathcal{Z} \geq 0$ and

$$\sup_{(n-1)t^{\frac{1}{2}} \leq |z| < nt^{\frac{1}{2}}} \omega_\alpha^{-2}(z, t) \lesssim n^{-2\alpha},$$

we get

$$\begin{aligned} & \mathbb{P}(\sup_{t>0} t^{-\frac{d}{2}} \sum_z \omega_\alpha^{-2}(z, t) \mathcal{Z}(\tau_z a) > \lambda) \\ & \leq \mathbb{P}(\exists t \in (0, +\infty), n \in \mathbb{N} : n^{-d} t^{-\frac{d}{2}} \sum_{(n-1)t^{\frac{1}{2}} \leq |z| < nt^{\frac{1}{2}}} \mathcal{Z}(\tau_z a) \gtrsim n^{2\alpha-d-2} \lambda) \\ & \leq \mathbb{P}(\exists t \in (0, +\infty), n \in \mathbb{N} : n^{-d} t^{-\frac{d}{2}} \sum_{|z| < nt^{\frac{1}{2}}} \mathcal{Z}(\tau_z a) \gtrsim n^{2\alpha-d-2} \lambda) \\ & \leq \sum_{m=1}^{\infty} \mathbb{P}(\exists t \in (0, +\infty), n \in \mathbb{N} : n^{-d} t^{-\frac{d}{2}} \sum_{|z| < nt^{\frac{1}{2}}} \mathcal{Z}(\tau_z a) \gtrsim m^{2\alpha-d-2} \lambda) \\ & \leq \sum_{m=1}^{\infty} \mathbb{P}(\sup_{R>0} R^{-d} \sum_{|z| < R} \mathcal{Z}(\tau_z a) \gtrsim m^{2\alpha-d-2} \lambda). \end{aligned}$$

We may now use in this last line the standard maximal function estimate ([1], Theorem 3.2)

$$\mathbb{P}(\sup_{R>0} R^{-d} \sum_{|z| < R} \mathcal{Z}(\tau_z a) > \delta) \leq \frac{\langle |\mathcal{Z}| \rangle}{\delta}$$

to conclude

$$\mathbb{P}(\sup_{t>0} t^{-\frac{d}{2}} \sum_z \omega_\alpha^{-2}(z, t) \mathcal{Z}(\tau_z a) > \lambda) \lesssim \sum_{m=1}^{\infty} \frac{\langle |\mathcal{Z}| \rangle}{m^{2\alpha-2-d} \lambda} \stackrel{\alpha > \frac{d}{2}+1}{\lesssim} \frac{\langle |\mathcal{Z}| \rangle}{\lambda}.$$

□

REFERENCES

- [1] M. A. Akcoglu and U. Krengel. Ergodic theorems for superadditive processes. *J. Reine Angew. Math.*, 323:53–67, 1981.
- [2] S. Armstrong, T. Kuusi, and J.-C. Mourrat. The additive structure of elliptic homogenization, preprint, arXiv:1602.00512.
- [3] S. N. Armstrong and C. K. Smart. Quantitative stochastic homogenization of convex integral functionals. *Ann. Sci. Éc. Norm. Supér.*, in press, arXiv:1406.0996.
- [4] P. de Buyer and J.-C. Mourrat. Diffusive decay of the environment viewed by the particle. *Electron. Commun. Probab.*, 20:no. 23, 12, 2015.
- [5] A. Gloria, S. Neukamm, and F. Otto. Quantification of ergodicity in stochastic homogenization: optimal bounds via spectral gap on Glauber dynamics. *Invent. Math.*, 199(2):455–515, 2015.
- [6] A. Gloria, S. Neukamm, and F. Otto. A regularity theory for random elliptic operators, preprint, arXiv:1409.2678.
- [7] A. Gloria and F. Otto. The corrector in stochastic homogenization: Near-optimal rates with optimal stochastic integrability, preprint, arXiv:1510.08290.
- [8] A. Lamacz, S. Neukamm, and F. Otto. Moment bounds for the corrector in stochastic homogenization of a percolation model. Preprint, arXiv:1406.5723, 2014.
- [9] J.-C. Mourrat. Variance decay for functionals of the environment viewed by the particle. *Ann. Inst. Henri Poincaré Probab. Stat.*, 47(1):294–327, 2011.
- [10] J.-C. Mourrat and F. Otto. Anchored nash inequalities and heat kernel bounds for static and dynamic degenerate environments. *J. Funct. Anal.*, in press, arXiv:1503.08280v1.

(Arianna Giunti) MAX PLANCK INSTITUTE FOR MATHEMATICS IN THE SCIENCES, LEIPZIG, GERMANY

(Jean-Christophe Mourrat) ECOLE NORMALE SUPÉRIEURE DE LYON, CNRS, LYON, FRANCE